

Contents

VII	GENERAL RELATIVITY	ii
23	From Special to General Relativity	1
23.1	Overview	1
23.2	Special Relativity Once Again	2
23.2.1	Geometric, Frame-Independent Formulation	2
23.2.2	Inertial Frames and Components of Vectors, Tensors and Physical Laws	4
23.2.3	Light Speed, the Interval, and Spacetime Diagrams	6
23.3	Differential Geometry in General Bases and in Curved Manifolds	8
23.3.1	Non-Orthonormal Bases	8
23.3.2	Vectors as Differential Operators; Tangent Space; Commutators	13
23.3.3	Differentiation of Vectors and Tensors; Connection Coefficients	15
23.3.4	Integration	21
23.4	The Stress-Energy Tensor Revisited	23
23.5	The Proper Reference Frame of an Accelerated Observer	27
23.5.1	Relation to Inertial Coordinates; Metric in Proper Reference Frame; Transport Law for Rotating Vectors	29
23.5.2	Geodesic Equation for Freely Falling Particle	31
23.5.3	Uniformly Accelerated Observer	32
23.5.4	Rindler Coordinates for Minkowski Spacetime	33

Part VII

GENERAL RELATIVITY

General Relativity

Version 1123.1.K.pdf, 25 April 2012

We have reached the final Part of this book, in which we present an introduction to the basic concepts of general relativity and its most important applications. This subject, although a little more challenging than the material that we have covered so far, is nowhere near as formidable as its reputation. Indeed, if you have mastered the techniques developed in the first five Parts, the path to the Einstein Field Equations should be short and direct.

The General Theory of Relativity is the crowning achievement of classical physics, the last great fundamental theory created prior to the discovery of quantum mechanics. Its formulation by Albert Einstein in 1915 marks the culmination of the great intellectual adventure undertaken by Newton 250 years earlier. Einstein created it after many wrong turns and with little experimental guidance, almost by pure thought. Unlike the special theory, whose physical foundations and logical consequences were clearly appreciated by physicists soon after Einstein's 1905 formulation, the unique and distinctive character of the general theory only came to be widely appreciated long after its creation. Ultimately, in hindsight, rival classical theories of gravitation came to seem unnatural, inelegant and arbitrary by comparison.¹

Experimental tests of Einstein's theory also were slow to come; only since 1970 have there been striking tests of high enough precision to convince most empiricists that, in all probability, and in its domain of applicability, general relativity is essentially correct. Despite this, it is still very poorly tested compared with, for example, quantum electrodynamics.

We begin our discussion of general relativity in Chap. 23 with a review, and an elaboration, of special relativity as developed in Chap. 2, focusing on those concepts that are crucial for the transition to general relativity. Our elaboration includes: (i) an extension of differential geometry to curvilinear coordinate systems and general bases both in the flat spacetime of special relativity and in the curved spacetime that is the venue for general relativity, (ii) an in-depth exploration of the stress-energy tensor, which in general relativity generates the curvature of spacetime, and (iii) construction and exploration of the reference frames of accelerated observers, e.g. physicists who reside on the Earth's surface. In Chap. 24, we turn to the basic concepts of general relativity, including spacetime curvature, the Einstein Field Equation that governs the generation of spacetime curvature, the laws of physics in curved spacetime, and weak-gravity limits of general relativity.

¹For a readable account at a popular level, see Will (1993); for a more detailed, scholarly account see, e.g. Pais (1982).

In the remaining chapters, we explore applications of general relativity to stars, black holes, gravitational waves, experimental tests of the theory, and cosmology. We begin in Chap 25 by studying the spacetime curvature around and inside highly compact stars (such as neutron stars). We then discuss the implosion of massive stars and describe the circumstances under which the implosion inevitably produces a black hole, we explore the surprising and, initially, counter-intuitive properties of black holes (both nonspinning holes and spinning holes), and we learn about the many-fingered nature of time in general relativity. In Chap. 26, we study experimental tests of general relativity, and then turn attention to gravitational waves, i.e. ripples in the curvature of spacetime that propagate with the speed of light. We explore the properties of these waves, their close analogy with electromagnetic waves, their production by binary stars and merging black holes, projects to detect them, both on earth and in space, and the prospects for using them to explore observationally the “warped side of the universe” and the nature of ultrastrong spacetime curvature. Finally, in Chap. 27 we draw upon all the previous Parts of this book, combining them with general relativity to describe the universe on the largest of scales and longest of times: cosmology. It is here, more than anywhere else in classical physics, that we are conscious of reaching a frontier where the still-promised land of quantum gravity beckons.

Chapter 23

From Special to General Relativity

Version 1123.1.K.pdf, 25 April 2012

Please send comments, suggestions, and errata via email to kip@caltech.edu or on paper to Kip Thorne, 350-17 Caltech, Pasadena CA 91125

Box 23.1 Reader's Guide

- This chapter relies significantly on
 - Chapter 2 on special relativity.
 - The discussion of connection coefficients in Sec. 11.5.
- This chapter is a foundation for the presentation of general relativity theory in Chaps. 24–27.

23.1 Overview

We begin our discussion of general relativity in this chapter with a review and elaboration of relevant material already covered in earlier chapters. In Sec. 23.2, we give a brief encapsulation of the special theory drawn largely from Chap. 2, emphasizing those aspects that underpin the transition to general relativity. Then in Sec. 23.3 we collect, review and extend the fundamental ideas of differential geometry that have been scattered throughout the book and which we shall need as foundations for the mathematics of *spacetime curvature* (Chap. 24); most importantly, we generalize differential geometry to encompass coordinate systems whose coordinate lines are not orthogonal and bases that are not orthonormal

Einstein's field equation (to be studied in Chap. 24) is a relationship between the curvature of spacetime and the matter that generates it, akin to the Maxwell equations' relationship between the electromagnetic field and the electric currents and charges that generate

it. The matter in Einstein’s equation is described by the *stress-energy tensor* that we introduced in Sec. 2.13. We revisit the stress-energy tensor in Sec. 23.4 and develop a deeper understanding of its properties.

In general relativity one often wishes to describe the outcome of measurements made by observers who refuse to fall freely—e.g., an observer who hovers in a spaceship just above the horizon of a black hole, or a gravitational-wave experimenter in an earth-bound laboratory. As a foundation for treating such observers, in Sec. 23.5 we examine measurements made by accelerated observers in the flat spacetime of special relativity.

23.2 Special Relativity Once Again

A pre-requisite to learning the theory of general relativity is to understand special relativity in geometric language. In Chap. 2, we discussed the foundations of special relativity with this in mind. In this section we briefly review the most important points.

We suggest that any reader who has not studied Chap. 2 read this Sec. 23.2 first, to get an overview and flavor of what will be important for our development of general relativity; and then (or in parallel with reading this Sec. 23.2), read those relevant sections of Chap. 2 that the reader does not already understand.

23.2.1 Geometric, Frame-Independent Formulation

In Secs. 1.1.1 and 2.2.2, we learned that *every law of physics must be expressible as a geometric, frame-independent relationship between geometric, frame-independent objects*. This is equally true in Newtonian physics, in special relativity and in general relativity. The key difference between the three is the geometric arena: In Newtonian physics the arena is 3-dimensional Euclidean space; in special relativity it is 4-dimensional Minkowski spacetime; in general relativity (Chap. 24) it is 4-dimensional curved spacetime; see Fig. 1 in the Introduction to Part I, and the associated discussion.

In special relativity, the demand that the laws be geometric relationships between geometric objects in Minkowski spacetime is called the *Principle of Relativity*; see Sec. 2.2.2. Examples of the geometric objects are: (i) A point \mathcal{P} in spacetime (which represents an *event*); Sec. 2.2.1. (ii) A parametrized curve in spacetime, such as the world line $\mathcal{P}(\tau)$ of a particle, for which the parameter τ is the particle’s *proper time*, i.e. the time measured by an ideal clock¹ that the particle carries (Fig. 23.1); Sec. 2.4.1. (iii) Vectors, such as the particle’s 4-velocity $\vec{u} = d\mathcal{P}/d\tau$ [the tangent vector to the curve $\mathcal{P}(\tau)$] and the particle’s 4-momentum $\vec{p} = m\vec{u}$ (with m the particle’s rest mass); Secs. 2.2.1 and 2.4.1. (iv) Tensors, such as the electromagnetic field tensor $\mathbf{F}(_, _)$; Secs. 1.3 and 2.3. A tensor, as we recall, is a linear real-valued function of vectors; when one puts vectors \vec{A} and \vec{B} into the slots of

¹Recall that an ideal clock is one that ticks uniformly when compared, e.g., to the period of the light emitted by some standard type of atom or molecule, and that has been made impervious to accelerations so two ideal clocks momentarily at rest with respect to each other tick at the same rate independent of their relative acceleration; see Secs. 2.2.1 and 2.4.1, and for greater detail, pp. 23–29 and 395–399 of Misner, Thorne and Wheeler (1973), henceforth cited as MTW.




Fig. 23.1: The world line $\mathcal{P}(\tau)$ of a particle in Minkowski spacetime and the tangent vector $\vec{u} = d\mathcal{P}/d\tau$ to this world line; \vec{u} is the particle's 4-velocity. The bending of the world line is produced by some force that acts on the particle, e.g. by the Lorentz force embodied in Eq. (23.3). Also shown is the light cone emitted from the event $\mathcal{P}(\tau = 1)$. Although the axes of an (arbitrary) inertial reference frame are shown, no reference frame is needed for the definition of the world line or its tangent vector \vec{u} or the light cone, or for the formulation of the Lorentz force law.

F, one obtains a real number (a scalar) $\mathbf{F}(\vec{A}, \vec{B})$ that is linear in \vec{A} and in \vec{B} so for example $\mathbf{F}(\vec{A}, b\vec{B} + c\vec{C}) = b\mathbf{F}(\vec{A}, \vec{B}) + c\mathbf{F}(\vec{A}, \vec{C})$. When one puts a vector \vec{B} into just one of the slots of **F** and leaves the other empty, one obtains a tensor with one empty slot, $\mathbf{F}(_, \vec{B})$, i.e. a vector. The result of putting a vector into the slot of a vector is the scalar product, $\vec{D}(\vec{B}) = \vec{D} \cdot \vec{B} = \mathbf{g}(\vec{D}, \vec{B})$, where $\mathbf{g}(_, _)$ is the metric.

In Secs. 2.3 and 2.4.1, we tied our definitions of the inner product and the spacetime metric to the ticking of ideal clocks: If $\Delta\vec{x}$ is the vector separation of two neighboring events $\mathcal{P}(\tau)$ and $\mathcal{P}(\tau + \Delta\tau)$ along a particle's world line, then

$$\boxed{\mathbf{g}(\Delta\vec{x}, \Delta\vec{x}) \equiv \Delta\vec{x} \cdot \Delta\vec{x} \equiv -(\Delta\tau)^2} . \quad (23.1)$$

This relation for any particle with any timelike world line, together with the linearity of $\mathbf{g}(_, _)$ in its two slots, is enough to determine **g** completely and to guarantee that it is symmetric, $\mathbf{g}(\vec{A}, \vec{B}) = \mathbf{g}(\vec{B}, \vec{A})$ for all \vec{A} and \vec{B} . Since the particle's 4-velocity \vec{u} is

$$\vec{u} = \frac{d\mathcal{P}}{d\tau} = \lim_{\Delta\tau \rightarrow 0} \frac{\mathcal{P}(\tau + \Delta\tau) - \mathcal{P}(\tau)}{\Delta\tau} \equiv \lim_{\Delta\tau \rightarrow 0} \frac{\Delta\vec{x}}{\Delta\tau} , \quad (23.2)$$

Eq. (23.1) implies that $\vec{u} \cdot \vec{u} = \mathbf{g}(\vec{u}, \vec{u}) = -1$ (Sec. 2.4.1).

The 4-velocity \vec{u} is an example of a *timelike* vector (Sec. 2.2.2); it has a negative inner product with itself (negative “squared length”). This shows up pictorially in the fact that \vec{u} lies inside the *light cone* (the cone swept out by the trajectories of photons emitted from the tail of \vec{u} ; see Fig. 23.1). Vectors \vec{k} on the light cone (the tangents to the world lines of the photons) are *null* and so have vanishing squared lengths, $\vec{k} \cdot \vec{k} = \mathbf{g}(\vec{k}, \vec{k}) = 0$; and vectors \vec{A} that lie outside the light cone are *spacelike* and have positive squared lengths, $\vec{A} \cdot \vec{A} > 0$. See Sec. 2.2.2.

An example of a physical law in 4-dimensional geometric language is the Lorentz force law (Sec. 2.4.2)

$$\frac{d\vec{p}}{d\tau} = q\mathbf{F}(_, \vec{u}) , \quad (23.3)$$

where q is the particle's charge and both sides of this equation are vectors, i.e. first-rank tensors, i.e. tensors with just one slot. As we learned in Secs. 1.5.1 and 2.5, it is convenient to give names to slots. When we do so, we can rewrite the Lorentz force law as

$$\frac{dp^\alpha}{d\tau} = qF^{\alpha\beta}u_\beta . \quad (23.4)$$

Here α is the name of the slot of the vector $dp^\alpha/d\tau$, α and β are the names of the slots of \mathbf{F} , β is the name of the slot of \mathbf{u} , and the double use of β with one up and one down on the right side of the equation represents the insertion of \vec{u} into the β slot of \mathbf{F} , whereby the two β slots disappear and we wind up with a vector whose slot is named α . As we learned in Sec. 1.5, this *slot-naming index notation* is isomorphic to the notation for components of vectors, tensors, and physical laws in some reference frame. However, no reference frames are needed or involved when one formulates the laws of physics in geometric, frame-independent language as above.

Those readers who do not feel completely comfortable with these concepts, statements and notation should reread the relevant portions of Chaps. 2 and 1.

EXERCISES

Exercise 23.1 *Practice: Frame-Independent Tensors*

Let \mathbf{A} , \mathbf{B} be second rank tensors.

- Show that $\mathbf{A} + \mathbf{B}$ is also a second rank tensor.
- Show that $\mathbf{A} \otimes \mathbf{B}$ is a fourth rank tensor.
- Show that the contraction of $\mathbf{A} \otimes \mathbf{B}$ on its first and fourth slots is a second rank tensor. (If necessary, consult Chap. 2 for a discussion of contraction).
- Write the following quantities in slot-naming index notation: the tensor $\mathbf{A} \otimes \mathbf{B}$; the simultaneous contraction of this tensor on its first and fourth slots and on its second and third slots.

23.2.2 Inertial Frames and Components of Vectors, Tensors and Physical Laws

In special relativity, a key role is played by *inertial reference frames*, Sec. 2.2.1. An inertial frame is an (imaginary) latticework of rods and clocks that moves through spacetime freely (inertially, without any force acting on it). The rods are orthogonal to each other and attached to inertial-guidance gyroscopes so they do not rotate. These rods are used to identify the spatial, Cartesian coordinates $(x^1, x^2, x^3) = (x, y, z)$ of an event \mathcal{P} [which we

also denote by lower case Latin indices $x^j(\mathcal{P})$ with j running over 1,2,3]. The latticework's clocks are ideal and are synchronized with each other via the Einstein light-pulse process. They are used to identify the temporal coordinate $x^0 = t$ of an event \mathcal{P} ; i.e. $x^0(\mathcal{P})$ is the time measured by that latticework clock whose world line passes through \mathcal{P} , at the moment of passage. The spacetime coordinates of \mathcal{P} are denoted by lower case Greek indices x^α , with α running over 0,1,2,3. An inertial frame's spacetime coordinates $x^\alpha(\mathcal{P})$ are called *Lorentz coordinates* or *inertial coordinates*.

In the real universe, spacetime curvature is very small in regions well-removed from concentrations of matter, e.g. in intergalactic space; so special relativity is highly accurate there. In such a region, frames of reference (rod-clock latticeworks) that are non-accelerating and non-rotating with respect to cosmologically distant galaxies (and thence with respect to a local frame in which the cosmic microwave radiation looks isotropic) constitute good approximations to inertial reference frames.

Associated with an inertial frame's Lorentz coordinates are basis vectors \vec{e}_α that point along the frame's coordinate axes (and thus are orthogonal to each other) and have unit length (making them orthonormal); Sec. 2.5. This *orthonormality* is embodied in the inner products

$$\boxed{\vec{e}_\alpha \cdot \vec{e}_\beta = \eta_{\alpha\beta}}, \quad (23.5)$$

where by definition

$$\boxed{\eta_{00} = -1, \quad \eta_{11} = \eta_{22} = \eta_{33} = +1, \quad \eta_{\alpha\beta} = 0 \text{ if } \alpha \neq \beta}. \quad (23.6)$$

Here and throughout Part VII (as in Chap. 2), we set the speed of light to unity (i.e. we use the *geometrized units* introduced in Sec. 1.10), so spatial lengths (e.g. along the x axis) and time intervals (e.g. along the t axis) are measured in the same units, seconds or meters with $1 \text{ s} = 2.99792458 \times 10^8 \text{ m}$.

In Sec. 2.5 (see also Sec. 1.5), we used the basis vectors of an inertial frame to build a component representation of tensor analysis. The fact that the inner products of timelike vectors with each other are negative, e.g. $\vec{e}_0 \cdot \vec{e}_0 = -1$, while those of spacelike vectors are positive, e.g. $\vec{e}_1 \cdot \vec{e}_1 = +1$, forced us to introduce two types of components: *covariant* (indices down) and *contravariant* (indices up). The covariant components of a tensor were computable by inserting the basis vectors into the tensor's slots, $u_\alpha = \vec{u}(\vec{e}_\alpha) \equiv \vec{u} \cdot \vec{e}_\alpha$; $F_{\alpha\beta} = \mathbf{F}(\vec{e}_\alpha, \vec{e}_\beta)$. For example, in our Lorentz basis the covariant components of the metric are $g_{\alpha\beta} = \mathbf{g}(\vec{e}_\alpha, \vec{e}_\beta) = \vec{e}_\alpha \cdot \vec{e}_\beta = \eta_{\alpha\beta}$. The contravariant components of a tensor were related to the covariant components via "index lowering" with the aid of the metric, $F_{\alpha\beta} = g_{\alpha\mu} g_{\beta\nu} F^{\mu\nu}$, which simply said that one reverses the sign when lowering a time index and makes no change of sign when lowering a space index. This lowering rule implied that the contravariant components of the metric in a Lorentz basis are the same numerically as the covariant components, $g^{\alpha\beta} = \eta_{\alpha\beta}$ and that they can be used to raise indices (i.e. to perform the trivial sign flip for temporal indices) $F^{\mu\nu} = g^{\mu\alpha} g^{\nu\beta} F_{\alpha\beta}$. As we saw in Sec. 2.5, tensors can be expressed in terms of their contravariant components as $\vec{p} = p^\alpha \vec{e}_\alpha$, and $\mathbf{F} = F^{\alpha\beta} \vec{e}_\alpha \otimes \vec{e}_\beta$, where \otimes represents the tensor product [Eqs. (1.5)].

We also learned in Chap. 2 that any frame independent geometric relation between tensors can be rewritten as a relation between those tensors' components in any chosen Lorentz

frame. When one does so, the resulting component equation takes *precisely the same form* as the slot-naming-index-notation version of the geometric relation (Sec. 1.5.1 and end of Sec. 2.5). For example, the component version of the Lorentz force law says $dp^\alpha/d\tau = qF^{\alpha\beta}u_\beta$, which is identical to Eq. (23.4). The only difference is the interpretation of the symbols. In the component equation $F^{\alpha\beta}$ are the components of \mathbf{F} and the repeated β in $F^{\alpha\beta}u_\beta$ is to be summed from 0 to 3. In the geometric relation $F^{\alpha\beta}$ means $\mathbf{F}(_, _)$ with the first slot named α and the second β , and the repeated β in $F^{\alpha\beta}u_\beta$ implies the insertion of \vec{u} into the second slot of \mathbf{F} to produce a single-slotted tensor, i.e. a vector whose slot is named α .

As we saw in Sec. 2.6, a particle's 4-velocity \vec{u} (defined originally without the aid of any reference frame; Fig. 23.1) has components, in any inertial frame, given by $u^0 = \gamma$, $u^j = \gamma v^j$ where $v^j = dx^j/dt$ is the particle's ordinary velocity and $\gamma \equiv 1/\sqrt{1 - \delta_{ij}v^i v^j}$. Similarly, the particle's energy $E \equiv p^0$ is $m\gamma$ and its spatial momentum is $p^j = m\gamma v^j$, i.e. in 3-dimensional geometric notation, $\mathbf{p} = m\gamma\mathbf{v}$. This is an example of the manner in which a choice of Lorentz frame produces a "3+1" split of the physics: a split of 4-dimensional spacetime into 3-dimensional space (with Cartesian coordinates x^j) plus 1-dimensional time $t = x^0$; a split of the particle's 4-momentum \vec{p} into its 3-dimensional spatial momentum \mathbf{p} and its 1-dimensional energy $E = p^0$; and similarly a split of the electromagnetic field tensor \mathbf{F} into the 3-dimensional electric field \mathbf{E} and 3-dimensional magnetic field \mathbf{B} ; cf. Secs. 2.6 and 2.11.

The principle of relativity (all laws expressible as geometric relations between geometric objects in Minkowski spacetime), when translated into 3+1 language, says that, *when the laws of physics are expressed in terms of components in a specific Lorentz frame, the form of those laws must be independent of one's choice of frame*. The components of tensors in one Lorentz frame are related to those in another by a Lorentz transformation (Sec. 2.7), so the principle of relativity can be restated as saying that, when expressed in terms of Lorentz-frame components, *the laws of physics must be Lorentz-invariant* (unchanged by Lorentz transformations). This is the version of the principle of relativity that one meets in most elementary treatments of special relativity. However, as the above discussion shows, it is a mere shadow of the true principle of relativity—the shadow cast onto Lorentz frames when one performs a 3+1 split. The ultimate, fundamental version of the principle of relativity is the one that needs no frames at all for its expression: *All the laws of physics are expressible as geometric relations between geometric objects that reside in Minkowski spacetime*.

23.2.3 Light Speed, the Interval, and Spacetime Diagrams

One set of physical laws that must be the same in all inertial frames is Maxwell's equations. Let us discuss the implications of Maxwell's equations for the speed of light c , momentarily abandoning geometrized units and returning to mks/SI units. According to Maxwell, c can be determined by performing non-radiative laboratory experiments; it is not necessary to measure the time it takes light to travel along some path. For example, measure the electrostatic force between two charges; that force is $\propto \epsilon_0^{-1}$, the electric permittivity of free space. Then allow one of these charges to travel down a wire and by its motion generate a magnetic field. Let the other charge move through this field and measure the magnetic force on it; that force is $\propto \mu_0$, the magnetic permittivity of free space. The ratio of these two forces

can be computed and is $\propto 1/\mu_0\epsilon_0 = c^2$. By combining the results of the two experiments, we therefore can deduce the speed of light c ; this is completely analogous to deducing the speed of seismic waves through rock from a knowledge of the rock's density and elastic moduli, using elasticity theory (Chap. 12). The principle of relativity, in operational form, dictates that the results of the electric and magnetic experiments must be independent of the Lorentz frame in which one chooses to perform them; therefore, the speed of light is frame-independent—as we argued by a different route in Sec. 2.2.2. It is this frame independence that enables us to introduce geometrized units with $c = 1$.

Another example of frame independence (Lorentz invariance) is provided by the *interval between two events* (Sec. 2.2.2). The components $g_{\alpha\beta} = \eta_{\alpha\beta}$ of the metric imply that, if $\Delta\vec{x}$ is the vector separating the two events and Δx^α are its components in some Lorentz coordinate system, then the squared length of $\Delta\vec{x}$ [also called the *interval* and denoted $(\Delta s)^2$] is given by

$$\boxed{(\Delta s)^2 \equiv \Delta\vec{x} \cdot \Delta\vec{x} = \mathbf{g}(\Delta\vec{x}, \Delta\vec{x}) = g_{\alpha\beta} \Delta x^\alpha \Delta x^\beta = -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2}. \quad (23.7)$$

Since $\Delta\vec{x}$ is a geometric, frame-independent object, so must be the interval. This implies that the equation $(\Delta s)^2 = -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2$ by which one computes the interval between the two chosen events in one Lorentz frame must give the same numerical result when used in any other frame; i.e., this expression must be Lorentz invariant. This *invariance of the interval* is the starting point for most introductions to special relativity—and, indeed, we used it as a starting point in Sec. 2.2.

Spacetime diagrams will play a major role in our development of general relativity. Accordingly, it is important that the reader feel very comfortable with them. We recommend reviewing Fig. 2.8 and Ex. 2.14.

EXERCISES

Exercise 23.2 *Example: Invariance of a Null Interval*

You have measured the intervals between a number of adjacent events in spacetime and thereby have deduced the metric \mathbf{g} . Your friend claims that the metric is some other frame-independent tensor $\tilde{\mathbf{g}}$ that differs from \mathbf{g} . Suppose that your correct metric \mathbf{g} and his wrong one $\tilde{\mathbf{g}}$ agree on the forms of the light cones in spacetime, i.e. they agree as to which intervals are null, which are spacelike and which are timelike; but they give different answers for the value of the interval in the spacelike and timelike cases, i.e. $\mathbf{g}(\Delta\vec{x}, \Delta\vec{x}) \neq \tilde{\mathbf{g}}(\Delta\vec{x}, \Delta\vec{x})$. Prove that $\tilde{\mathbf{g}}$ and \mathbf{g} differ solely by a scalar multiplicative factor, $\tilde{\mathbf{g}} = a\mathbf{g}$ for some scalar a . [*Hint*: pick some Lorentz frame and perform computations there, then lift yourself back up to a frame-independent viewpoint.]

Exercise 23.3 *Problem: Causality*

If two events occur at the same spatial point but not simultaneously in one inertial frame, prove that the temporal order of these events is the same in all inertial frames. Prove also

that in all other frames the temporal interval Δt between the two events is larger than in the first frame, and that there are no limits on the events' spatial or temporal separation in the other frames. Give *two* proofs of these results, one algebraic and the other via spacetime diagrams.

23.3 Differential Geometry in General Bases and in Curved Manifolds

The differential geometry (tensor-analysis) formalism reviewed in the last section is inadequate for general relativity in several ways:

First, in general relativity we shall need to use bases \vec{e}_α that are not orthonormal, i.e. for which $\vec{e}_\alpha \cdot \vec{e}_\beta \neq \eta_{\alpha\beta}$. For example, near a spinning black hole there is much power in using a time basis vector \vec{e}_t that is tied in a simple way to the metric's time-translation symmetry and a spatial basis vector \vec{e}_ϕ that is tied to its rotational symmetry. This time basis vector has an inner product with itself $\vec{e}_t \cdot \vec{e}_t = g_{tt}$ that is influenced by the slowing of time near the hole so $g_{tt} \neq -1$; and \vec{e}_ϕ is not orthogonal to \vec{e}_t , $\vec{e}_t \cdot \vec{e}_\phi = g_{t\phi} \neq 0$, as a result of the dragging of inertial frames by the hole's spin. In this section, we shall generalize our formalism to treat such non-orthonormal bases.

Second, in the curved spacetime of general relativity (and in any other curved space, e.g. the two-dimensional surface of the earth), the definition of a vector as an arrow connecting two points (Secs. 1.2 and 2.2.1) is suspect, as it is not obvious on what route the arrow should travel nor that the linear algebra of tensor analysis should be valid for such arrows. In this section we shall refine the concept of a vector to deal with this problem, and in the process we shall find ourselves introducing the concept of a *tangent space* in which the linear algebra of tensors takes place—a different tangent space for tensors that live at different points in the space.

Third, once we have been forced to think of a tensor as residing in a specific tangent space at a specific point in the space, there arises the question of how one can transport tensors from the tangent space at one point to the tangent space at an adjacent point. Since the notion of a gradient of a vector depends on comparing the vector at two different points and thus depends on the details of transport, we will have to rework the notion of a gradient and the gradient's connection coefficients; and since, in doing an integral, one must add contributions that live at different points in the space, we must also rework the notion of integration.

We shall tackle each of these three issues in turn in the following four subsections.

23.3.1 Non-Orthonormal Bases

Consider an n -dimensional *manifold*, i.e., a space that, in the neighborhood of any point, has the same topological and smoothness properties as n -dimensional Euclidean space, though




Fig. 23.2: Non-orthonormal basis vectors \vec{e}_j in Euclidean 3-space and two members \vec{e}^1 and \vec{e}^3 of the dual basis. The vectors \vec{e}_1 and \vec{e}_2 lie in the horizontal plane, so \vec{e}^3 is orthogonal to that plane, i.e. it points vertically upward, and its inner product with \vec{e}_3 is unity. Similarly, the vectors \vec{e}_2 and \vec{e}_3 span a vertical plane, so \vec{e}^1 is orthogonal to that plane, i.e. it points horizontally, and its inner product with \vec{e}_1 is unity.

it might *not* have a locally Euclidean metric and perhaps no metric at all. Examples that do have metrics are 4-dimensional spacetime, 3-dimensional Euclidean space, and the 2-dimensional surface of a sphere. In this chapter, all manifolds we consider will have metrics.

At some point \mathcal{P} in our chosen n -dimensional manifold with metric, introduce a set of basis vectors $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ and denote them generally as \vec{e}_α . We seek to generalize the formalism of Sec. 23.2 in such a way that the index manipulation rules for components of tensors are unchanged. For example, we still want it to be true that covariant components of any tensor are computable by inserting the basis vectors into the tensor's slots, $F_{\alpha\beta} = \mathbf{F}(\vec{e}_\alpha, \vec{e}_\beta)$, and that the tensor itself can be reconstructed from its contravariant components as $\mathbf{F} = F^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu$, and that the two sets of components are computable from each other via raising and lowering with the metric components, $F_{\alpha\beta} = g_{\alpha\mu} g_{\beta\nu} F^{\mu\nu}$. The only thing we do not want to preserve is the orthonormal values of the metric components; i.e. we must allow the basis to be nonorthonormal and thus $\vec{e}_\alpha \cdot \vec{e}_\beta = g_{\alpha\beta}$ to have arbitrary values (except that the metric should be nondegenerate, so no linear combination of the \vec{e}_α 's vanishes, which means that the matrix $\|g_{\alpha\beta}\|$ should have nonzero determinant).

We can easily achieve our goal by introducing a second set of basis vectors, denoted $\{\vec{e}^1, \vec{e}^2, \dots, \vec{e}^n\}$, which is *dual* to our first set in the sense that

$$\boxed{\vec{e}^\mu \cdot \vec{e}_\beta \equiv \mathbf{g}(\vec{e}^\mu, \vec{e}_\beta) = \delta^\mu_\beta}. \quad (23.8)$$

Here δ^α_β is the Kronecker delta. This duality relation actually constitutes a *definition* of the e^μ once the \vec{e}_α have been chosen. To see this, regard \vec{e}^μ as a tensor of rank one. This tensor is defined as soon as its value on each and every vector has been determined. Expression (23.8) gives the value $\vec{e}^\mu(\vec{e}_\beta) = \vec{e}^\mu \cdot \vec{e}_\beta$ of \vec{e}^μ on each of the four basis vectors \vec{e}_β ; and since every other vector can be expanded in terms of the \vec{e}_β 's and $\vec{e}^\mu(_)$ is a linear function, Eq. (23.8) thereby determines the value of \vec{e}^μ on every other vector.

The duality relation (23.8) says that \vec{e}^1 is always perpendicular to all the \vec{e}_α except \vec{e}_1 ; and its scalar product with \vec{e}_1 is unity—and similarly for the other basis vectors. This interpretation is illustrated for 3-dimensional Euclidean space in Fig. 23.2. In Minkowski spacetime, if \vec{e}_α are an orthonormal Lorentz basis, then duality dictates that $\vec{e}^0 = -\vec{e}_0$, and $\vec{e}^j = +\vec{e}_j$.

The duality relation (23.8) leads immediately to the same index-manipulation formalism as we have been using, if one defines the contravariant, covariant and mixed components of tensors in the obvious manner

$$\boxed{F^{\mu\nu} = \mathbf{F}(\vec{e}^\mu, \vec{e}^\nu), \quad F_{\alpha\beta} = \mathbf{F}(\vec{e}_\alpha, \vec{e}_\beta), \quad F^\mu{}_\beta = \mathbf{F}(\vec{e}^\mu, \vec{e}_\beta)}; \quad (23.9)$$

see Ex. 23.4. Among the consequences of this duality are the following: (i)

$$\boxed{g^{\mu\beta} g_{\nu\beta} = \delta_\nu^\mu}, \quad (23.10)$$

i.e., the matrix of contravariant components of the metric is inverse to that of the covariant components, $\|g^{\mu\nu}\| = \|g_{\alpha\beta}\|^{-1}$; this relation guarantees that when one raises indices on a tensor $F_{\alpha\beta}$ with $g^{\mu\alpha}$ and then lowers them back down with $g_{\nu\beta}$, one recovers one's original covariant components $F_{\alpha\beta}$ unaltered. (ii)

$$\boxed{\mathbf{F} = F^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu = F_{\alpha\beta} \vec{e}^\alpha \otimes \vec{e}^\beta = F^\mu{}_\beta \vec{e}_\mu \otimes \vec{e}^\beta}, \quad (23.11)$$

i.e., one can reconstruct a tensor from its components by lining up the indices in a manner that accords with the rules of index manipulation. (iii)

$$\boxed{\mathbf{F}(\vec{p}, \vec{q}) = F^{\alpha\beta} p_\alpha p_\beta}, \quad (23.12)$$

i.e., the component versions of tensorial equations are identical in mathematical symbology to the slot-naming-index-notation versions.

Associated with any coordinate system $x^\alpha(\mathcal{P})$ there is a *coordinate basis* whose basis vectors are defined by

$$\boxed{\vec{e}_\alpha \equiv \frac{\partial \mathcal{P}}{\partial x^\alpha}}. \quad (23.13)$$

Since the derivative is taken holding the other coordinates fixed, the basis vector \vec{e}_α points along the α coordinate axis (the axis on which x^α changes and all the other coordinates are held fixed).

In an orthogonal curvilinear coordinate system, e.g. circular polar coordinates (ϖ, ϕ) in Euclidean 2-space, this coordinate basis is quite different from the coordinate system's orthonormal basis. For example, $\vec{e}_\phi = (\partial \mathcal{P} / \partial \phi)_\varpi$ is a very long vector at large radii and a very short vector at small radii (cf. Fig. 23.3); the corresponding unit-length vector is $\vec{e}_\phi^\hat{=} = (1/\varpi)\vec{e}_\phi$. By contrast, $\vec{e}_\varpi = (\partial \mathcal{P} / \partial \varpi)_\phi$ already has unit length, so the corresponding orthonormal basis vector is simply $\vec{e}_\varpi^\hat{=} = \vec{e}_\varpi$. The metric components in the coordinate basis are readily seen to be $g_{\phi\phi} = \varpi^2$, $g_{\varpi\varpi} = 1$, $g_{\varpi\phi} = g_{\phi\varpi} = 0$, which are in accord with the equation for the squared distance (interval) between adjacent points $ds^2 = g_{ij} dx^i dx^j = d\varpi^2 + \varpi^2 d\phi^2$. The metric components in the orthonormal basis, of course, are $g_{i\hat{j}} = \delta_{ij}$.




Fig. 23.3: A circular coordinate system $\{\varpi, \phi\}$ and its coordinate basis vectors $\vec{e}_\varpi = \partial\mathcal{P}/\partial\varpi$, $\vec{e}_\phi = \partial\mathcal{P}/\partial\phi$ at several locations in the coordinate system. Also shown is the orthonormal basis vector \hat{e}_ϕ .

Henceforth, we shall use hats to identify orthonormal bases; bases whose indices do not have hats will typically (though not always) be coordinate bases.

In general, we can construct the basis $\{\vec{e}^\mu\}$ that is dual to the coordinate basis $\{\vec{e}_\alpha\} = \{\partial\mathcal{P}/\partial x^\alpha\}$ by taking the gradients of the coordinates, viewed as scalar fields $x^\alpha(\mathcal{P})$:

$$\boxed{\vec{e}^\mu = \vec{\nabla}x^\mu} . \quad (23.14)$$

It is straightforward to verify the duality relation (23.8) for these two bases:

$$\vec{e}^\mu \cdot \vec{e}_\alpha = \vec{e}_\alpha \cdot \vec{\nabla}x^\mu = \nabla_{\vec{e}_\alpha}x^\mu = \nabla_{\partial\mathcal{P}/\partial x^\alpha}x^\mu = \frac{\partial x^\mu}{\partial x^\alpha} = \delta_\alpha^\mu . \quad (23.15)$$

In any coordinate system, the expansion of the metric in terms of the dual basis, $\mathbf{g} = g_{\alpha\beta}\vec{e}^\alpha \otimes \vec{e}^\beta = g_{\alpha\beta}\vec{\nabla}x^\alpha \otimes \vec{\nabla}x^\beta$ is intimately related to the line element $ds^2 = g_{\alpha\beta}dx^\alpha dx^\beta$: Consider an infinitesimal vectorial displacement $d\vec{x} = dx^\alpha(\partial/\partial x^\alpha)$. Insert this displacement into the metric's two slots, to obtain the interval ds^2 along $d\vec{x}$. The result is $ds^2 = g_{\alpha\beta}\nabla x^\alpha \otimes \nabla x^\beta(d\vec{x}, d\vec{x}) = g_{\alpha\beta}(d\vec{x} \cdot \nabla x^\alpha)(d\vec{x} \cdot \nabla x^\beta) = g_{\alpha\beta}dx^\alpha dx^\beta$; i.e.

$$\boxed{ds^2 = g_{\alpha\beta}dx^\alpha dx^\beta} . \quad (23.16)$$

Here the second equality follows from the definition of the tensor product \otimes , and the third from the fact that for any scalar field ψ , $d\vec{x} \cdot \nabla\psi$ is the change $d\psi$ along $d\vec{x}$.

Any two bases $\{\vec{e}_\alpha\}$ and $\{\vec{e}_\mu\}$ can be expanded in terms of each other:

$$\boxed{\vec{e}_\alpha = \vec{e}_\mu L^\mu{}_\alpha, \quad \vec{e}_\mu = \vec{e}_\alpha L^\alpha{}_\mu} . \quad (23.17)$$

(Note: by convention the first index on L is always placed up and the second is always placed down.) The quantities $\|L^\mu{}_\alpha\|$ and $\|L^\alpha{}_\mu\|$ are transformation matrices and since they operate in opposite directions, they must be the inverse of each other

$$\boxed{L^\mu{}_\alpha L^\alpha{}_\nu = \delta^\mu{}_\nu, \quad L^\alpha{}_\mu L^\mu{}_\beta = \delta^\alpha{}_\beta} . \quad (23.18)$$

These $\|L^{\bar{\mu}}_{\alpha}\|$ are the generalizations of Lorentz transformations to arbitrary bases; cf. Eqs. (2.35a), (2.35b). As in the Lorentz-transformation case, the transformation laws (23.17) for the basis vectors imply corresponding transformation laws for components of vectors and tensors—laws that entail lining up indices in the obvious manner; e.g.

$$\boxed{A_{\bar{\mu}} = L^{\alpha}_{\bar{\mu}} A_{\alpha}, \quad T^{\bar{\mu}\bar{\nu}}_{\bar{\rho}} = L^{\bar{\mu}}_{\alpha} L^{\bar{\nu}}_{\beta} L^{\gamma}_{\bar{\rho}} T^{\alpha\beta}_{\gamma}, \quad \text{and similarly in the opposite direction.}} \quad (23.19)$$

For coordinate bases, these $L^{\bar{\mu}}_{\alpha}$ are simply the partial derivatives of one set of coordinates with respect to the other

$$\boxed{L^{\bar{\mu}}_{\alpha} = \frac{\partial x^{\bar{\mu}}}{\partial x^{\alpha}}, \quad L^{\alpha}_{\bar{\mu}} = \frac{\partial x^{\alpha}}{\partial x^{\bar{\mu}}},} \quad (23.20)$$

as one can easily deduce via

$$\vec{e}_{\alpha} = \frac{\partial \mathcal{P}}{\partial x^{\alpha}} = \frac{\partial x^{\mu}}{\partial x^{\alpha}} \frac{\partial \mathcal{P}}{\partial x^{\mu}} = \vec{e}_{\mu} \frac{\partial x^{\mu}}{\partial x^{\alpha}}. \quad (23.21)$$

In many physics textbooks a tensor is *defined* as a set of components $F_{\alpha\beta}$ that obey the transformation laws

$$F_{\alpha\beta} = F_{\mu\nu} \frac{\partial x^{\mu}}{\partial x^{\alpha}} \frac{\partial x^{\nu}}{\partial x^{\beta}}. \quad (23.22)$$

This definition is in accord with Eqs. (23.19) and (23.20), though it hides the true and very simple nature of a tensor as a linear function of frame-independent vectors.

EXERCISES

Exercise 23.4 *Derivation: Index Manipulation Rules from Duality*

For an arbitrary basis $\{\vec{e}_{\alpha}\}$ and its dual basis $\{\vec{e}^{\mu}\}$, use (i) the duality relation (23.8), (ii) the definition (23.9) of components of a tensor, and (iii) the relation $\vec{A} \cdot \vec{B} = \mathbf{g}(\vec{A}, \vec{B})$ between the metric and the inner product to deduce the following results:

- (a) The relations

$$\vec{e}^{\mu} = g^{\mu\alpha} \vec{e}_{\alpha}, \quad \vec{e}_{\alpha} = g_{\alpha\mu} \vec{e}^{\mu}. \quad (23.23)$$

- (b) The fact that indices on the components of tensors can be raised and lowered using the components of the metric, e.g.

$$F^{\mu\nu} = g^{\mu\alpha} F_{\alpha}{}^{\nu}, \quad p_{\alpha} = g_{\alpha\beta} p^{\beta}. \quad (23.24)$$

- (c) The fact that a tensor can be reconstructed from its components in the manner of Eq. (23.11).

Exercise 23.5 *Practice: Transformation Matrices for Circular Polar Bases*

Consider the circular coordinate system $\{\varpi, \phi\}$ and its coordinate bases and orthonormal bases as discussed in Fig. 23.3 and the associated text. These coordinates are related to Cartesian coordinates $\{x, y\}$ by the usual relations $x = \varpi \cos \phi$, $y = \varpi \sin \phi$.

- (a) Evaluate the components (L^x_{ϖ} etc.) of the transformation matrix that links the two coordinate bases $\{\vec{e}_x, \vec{e}_y\}$ and $\{\vec{e}_{\varpi}, \vec{e}_{\phi}\}$. Also evaluate the components (L^{ϖ}_x etc.) of the inverse transformation matrix.
- (b) Evaluate, similarly, the components of the transformation matrix and its inverse linking the bases $\{\vec{e}_x, \vec{e}_y\}$ and $\{\vec{e}_{\hat{\varpi}}, \vec{e}_{\hat{\phi}}\}$.
- (c) Consider the vector $\vec{A} \equiv \vec{e}_x + 2\vec{e}_y$. What are its components in the other two bases?

23.3.2 Vectors as Differential Operators; Tangent Space; Commutators

As was discussed above, the notion of a vector as an arrow connecting two points is problematic in a curved manifold, and must be refined. As a first step in the refinement, let us consider the tangent vector \vec{A} to a curve $\mathcal{P}(\zeta)$ at some point $\mathcal{P}_o \equiv \mathcal{P}(\zeta = 0)$. We have defined that tangent vector by the limiting process

$$\vec{A} \equiv \frac{d\mathcal{P}}{d\zeta} \equiv \lim_{\Delta\zeta \rightarrow 0} \frac{\mathcal{P}(\Delta\zeta) - \mathcal{P}(0)}{\Delta\zeta} \quad (23.25)$$

[Eq. (23.2)]. In this definition the difference $\mathcal{P}(\zeta) - \mathcal{P}(0)$ means the tiny arrow reaching from $\mathcal{P}(0) \equiv \mathcal{P}_o$ to $\mathcal{P}(\Delta\zeta)$. In the limit as $\Delta\zeta$ becomes vanishingly small, these two points get arbitrarily close together; and in such an arbitrarily small region of the manifold, the effects of the manifold's curvature become arbitrarily small and negligible (just think of an arbitrarily tiny region on the surface of a sphere), so the notion of the arrow should become sensible. However, before the limit is completed, we are required to divide by $\Delta\zeta$, which makes our arbitrarily tiny arrow big again. What meaning can we give to this?

One way to think about it is to imagine embedding the curved manifold in a higher dimensional flat space (e.g., embed the surface of a sphere in a flat 3-dimensional Euclidean space as shown in Fig. 23.4). Then the tiny arrow $\mathcal{P}(\Delta\zeta) - \mathcal{P}(0)$ can be thought of equally well as lying on the sphere, or as lying in a surface that is tangent to the sphere and is flat, as measured in the flat embedding space. We can give meaning to $[\mathcal{P}(\Delta\zeta) - \mathcal{P}(0)]/\Delta\zeta$ if we regard this as a formula for lengthening an arrow-type vector in the flat tangent surface; correspondingly, we must regard the resulting tangent vector \vec{A} as an arrow living in the tangent surface.

The (conceptual) flat tangent surface at the point \mathcal{P}_o is called the *tangent space* to the curved manifold at that point. It has the same number of dimensions n as the manifold itself (two in the case of Fig. 23.4). Vectors at \mathcal{P}_o are arrows residing in that point's tangent space, tensors at \mathcal{P}_o are linear functions of these vectors, and all the linear algebra of vectors and tensors that reside at \mathcal{P}_o occurs in this tangent space. For example, the inner product of two vectors \vec{A} and \vec{B} at \mathcal{P}_o (two arrows living in the tangent space there) is computed via

the standard relation $\vec{A} \cdot \vec{B} = \mathbf{g}(\vec{A}, \vec{B})$ using the metric \mathbf{g} that also resides in the tangent space.

This pictorial way of thinking about the tangent space and vectors and tensors that reside in it is far too heuristic to satisfy most mathematicians. Therefore, mathematicians have insisted on making it much more precise at the price of greater abstraction: *Mathematicians define the tangent vector to the curve $\mathcal{P}(\zeta)$ to be the derivative $d/d\zeta$ which differentiates scalar fields along the curve.* This derivative operator is very well defined by the rules of ordinary differentiation; if $\psi(\mathcal{P})$ is a scalar field in the manifold, then $\psi[\mathcal{P}(\zeta)]$ is a function of the real variable ζ , and its derivative $(d/d\zeta)\psi[\mathcal{P}(\zeta)]$ evaluated at $\zeta = 0$ is the ordinary derivative of elementary calculus. Since the derivative operator $d/d\zeta$ differentiates in the manifold along the direction in which the curve is moving, it is often called the *directional derivative* along $\mathcal{P}(\zeta)$. Mathematicians notice that all the directional derivatives at a point \mathcal{P}_o of the manifold form a vector space (they can be multiplied by scalars and added and subtracted to get new vectors), and so they define this vector space to be the tangent space at \mathcal{P}_o .

This mathematical procedure turns out to be isomorphic to the physicists' more heuristic way of thinking about the tangent space. In physicists' language, if one introduces a coordinate system in a region of the manifold containing \mathcal{P}_o and constructs the corresponding coordinate basis $\vec{e}_\alpha = \partial\mathcal{P}/\partial x^\alpha$, then one can expand any vector in the tangent space as $\vec{A} = A^\alpha \partial\mathcal{P}/\partial x^\alpha$. One can also construct, in physicists' language, the directional derivative along \vec{A} ; it is $\partial_{\vec{A}} \equiv A^\alpha \partial/\partial x^\alpha$. Evidently, the components A^α of the physicist's vector \vec{A} (an arrow) are identical to the coefficients A^α in the coordinate-expansion of the directional derivative $\partial_{\vec{A}}$. There therefore is a one-to-one correspondence between the directional derivatives $\partial_{\vec{A}}$ at \mathcal{P}_o and the vectors \vec{A} there, and a complete isomorphism between the tangent-space manipulations that a mathematician will perform treating the directional derivatives as vectors, and those that a physicist will perform treating the arrows as vectors.




Fig. 23.4: A curve $\mathcal{P}(\zeta)$ on the surface of a sphere and the curve's tangent vector $\vec{A} = d\mathcal{P}/d\zeta$ at $\mathcal{P}(\zeta = 0) \equiv \mathcal{P}_o$. The tangent vector lives in the tangent space at \mathcal{P}_o , i.e. in the flat plane that is tangent to the sphere there as seen in the flat Euclidean 3-space in which the sphere's surface is embedded.

“Why not abandon the fuzzy concept of a vector as an arrow, and *redefine the vector \vec{A} to be the same as the directional derivative $\partial_{\vec{A}}$?*” mathematicians have demanded of physicists. Slowly, over the past century, physicists have come to see the merit in this approach: (i) It does, indeed, make the concept of a vector more rigorous than before. (ii) It simplifies a number of other concepts in mathematical physics, e.g., the commutator of two vector fields; see below. (iii) It facilitates communication with mathematicians. With these motivations in mind, and because one always gains conceptual and computational power by having multiple viewpoints at one’s finger tips (see, e.g., Feynman, 1966), we shall regard vectors henceforth *both* as arrows living in a tangent space and as directional derivatives. Correspondingly, we shall assert the equalities

$$\boxed{\frac{\partial \mathcal{P}}{\partial x^\alpha} = \frac{\partial}{\partial x^\alpha} \quad , \quad \vec{A} = \partial_{\vec{A}}} \quad , \quad (23.26)$$

and shall often expand vectors in a coordinate basis using the notation

$$\boxed{\vec{A} = A^\alpha \frac{\partial}{\partial x^\alpha}} \quad . \quad (23.27)$$

This directional-derivative viewpoint on vectors makes natural the concept of the *commutator* of two vector fields \vec{A} and \vec{B} : $[\vec{A}, \vec{B}]$ is the vector which, when viewed as a differential operator, is given by $[\partial_{\vec{A}}, \partial_{\vec{B}}]$ —where the latter quantity is the same commutator as one meets elsewhere in physics, e.g. in quantum mechanics. Using this definition, we can compute the components of the commutator in a coordinate basis:

$$\boxed{[\vec{A}, \vec{B}] \equiv \left[A^\alpha \frac{\partial}{\partial x^\alpha}, B^\beta \frac{\partial}{\partial x^\beta} \right] = \left(A^\alpha \frac{\partial B^\beta}{\partial x^\alpha} - B^\alpha \frac{\partial A^\beta}{\partial x^\alpha} \right) \frac{\partial}{\partial x^\beta}} \quad . \quad (23.28)$$

This is an operator equation where the final derivative is presumed to operate on a scalar field just as in quantum mechanics. From this equation we can read off the components of the commutator in any coordinate basis; they are $A^\alpha B^\beta_{,\alpha} - B^\alpha A^\beta_{,\alpha}$, where the comma denotes partial differentiation. Figure 23.5 uses this equation to deduce the geometric meaning of the commutator: it is the fifth leg needed to close a quadrilateral whose other four legs are constructed from the vector fields \vec{A} and \vec{B} .

The commutator is useful as a tool for distinguishing between coordinate bases and non-coordinate bases (also called non-holonomic bases): In a coordinate basis, the basis vectors are just the coordinate system’s partial derivatives, $\vec{e}_\alpha = \partial/\partial x^\alpha$, and since partial derivatives commute, it must be that $[\vec{e}_\alpha, \vec{e}_\beta] = 0$. Conversely (as Fig. 23.5 explains), if one has a basis with vanishing commutators $[\vec{e}_\alpha, \vec{e}_\beta] = 0$, then it is possible to construct a coordinate system for which this is the coordinate basis. In a non-coordinate basis, at least one of the commutators $[\vec{e}_\alpha, \vec{e}_\beta]$ will be nonzero.

23.3.3 Differentiation of Vectors and Tensors; Connection Coefficients

In a curved manifold, the differentiation of vectors and tensors is rather subtle. To elucidate the problem, let us recall how we defined such differentiation in Minkowski spacetime or

Euclidean space (Sec. 1.7). Converting to the above notation, we began by defining the directional derivative of a tensor field $\mathbf{F}(\mathcal{P})$ along the tangent vector $\vec{A} = d/d\zeta$ to a curve $\mathcal{P}(\zeta)$:

$$\nabla_{\vec{A}}\mathbf{F} \equiv \lim_{\Delta\zeta \rightarrow 0} \frac{\mathbf{F}[\mathcal{P}(\Delta\zeta)] - \mathbf{F}[\mathcal{P}(0)]}{\Delta\zeta}. \quad (23.29)$$

This definition is problematic because $\mathbf{F}[\mathcal{P}(\Delta\zeta)]$ lives in a different tangent space than $\mathbf{F}[\mathcal{P}(0)]$. To make the definition meaningful, we must identify some *connection* between the two tangent spaces, when their points $\mathcal{P}(\Delta\zeta)$ and $\mathcal{P}(0)$ are arbitrarily close together. That connection is equivalent to identifying a rule for transporting \mathbf{F} from one tangent space to the other.

In flat space or flat spacetime, and when \mathbf{F} is a vector \vec{F} , that transport rule is obvious: keep \vec{F} parallel to itself and keep its length fixed during the transport; in other words, keep constant its components in an orthonormal coordinate system (Cartesian coordinates in Euclidean space, Lorentz coordinates in Minkowski spacetime). This is called the *law of parallel transport*. For a tensor \mathbf{F} , the parallel transport law is the same: keep its components fixed in an orthonormal coordinate basis.

Now, just as the curvature of the earth's surface prevents one from placing a Cartesian coordinate system on it, so nonzero curvature of any other manifold prevents one from introducing orthonormal coordinates; see Chap. 24. However, in an arbitrarily small region on the earth's surface, one can introduce coordinates that are arbitrarily close to Cartesian (as surveyors well know); the fractional deviations from Cartesian need be no larger than




Fig. 23.5: The commutator $[\vec{A}, \vec{B}]$ of two vector fields. In this diagram the vectors are assumed to be so small that the curvature of the manifold is negligible in the region of the diagram, so all the vectors can be drawn lying in the surface itself rather than in their respective tangent spaces. In evaluating the two terms in the commutator (23.28), a locally orthonormal coordinate basis is used, so $A^\alpha \partial B^\beta / \partial x^\alpha$ is the amount by which the vector \vec{B} changes when one travels along \vec{A} (i.e. it is the rightward & downward pointing dashed curve in the upper right), and $B^\alpha \partial A^\beta / \partial x^\alpha$ is the amount by which \vec{A} changes when one travels along \vec{B} (i.e. it is the rightward & upward pointing dashed curve). According to Eq. (23.28), the difference of these two dashed curves is the commutator $[\vec{A}, \vec{B}]$. As the diagram shows, this commutator closes the quadrilateral whose legs are \vec{A} and \vec{B} . If the commutator vanishes, then there is no gap in the quadrilateral, which means that in the region covered by this diagram one can construct a coordinate system in which \vec{A} and \vec{B} are coordinate basis vectors.

$\mathcal{O}(L^2/R^2)$, where L is the size of the region and R is the earth's radius (see Sec. 24.3). Similarly, in curved spacetime, in an arbitrarily small region, one can introduce coordinates that are arbitrarily close to Lorentz, differing only by amounts quadratic in the size of the region — and similarly for a *local* orthonormal coordinate basis in any curved manifold.

When defining $\nabla_{\vec{A}}\mathbf{F}$, one is sensitive only to first order changes of quantities, not second, so the parallel transport used in defining it in a flat manifold, based on constancy of components in an orthonormal coordinate basis, must also work in a *local* orthonormal coordinate basis of any curved manifold: In Eq. (23.29) one must transport \mathbf{F} from $\mathcal{P}(\Delta\zeta)$ to $\mathcal{P}(0)$, holding its components fixed in a locally orthonormal coordinate basis (parallel transport), and then take the difference in the tangent space at $\mathcal{P}_o = \mathcal{P}(0)$, divide by $\Delta\zeta$, and let $\Delta\zeta \rightarrow 0$. The result is a tensor at \mathcal{P}_o : the directional derivative $\nabla_{\vec{A}}\mathbf{F}$ of \mathbf{F} .

Having made the directional derivative meaningful, one can proceed as in Secs. 1.7 and 2.10: define the gradient of \mathbf{F} by $\nabla_{\vec{A}}\mathbf{F} = \vec{\nabla}\mathbf{F}(_, _, \vec{A})$ [i.e., put \vec{A} in the last, differentiation, slot of $\vec{\nabla}\mathbf{F}$; Eq. (1.15b)].

As in Chap. 2, in any basis we denote the components of $\vec{\nabla}\mathbf{F}$ by $F_{\alpha\beta;\gamma}$; and as in Sec. 11.5 (elasticity theory), we can compute these components in any basis with the aid of that basis's *connection coefficients*.

In Sec. 11.5, we restricted ourselves to an orthonormal basis in Euclidean space and thus had no need to distinguish between covariant and contravariant indices; all indices were written as subscripts. Now, with non-orthonormal bases and in spacetime, we must distinguish covariant and contravariant indices. Accordingly, by analogy with Eq. (11.38), we define the connection coefficients $\Gamma^\mu_{\alpha\beta}$ as

$$\boxed{\nabla_\beta \vec{e}_\alpha \equiv \nabla_{\vec{e}_\beta} \vec{e}_\alpha = \Gamma^\mu_{\alpha\beta} \vec{e}_\mu}, \quad (23.30)$$

where $\nabla_\beta \equiv \nabla_{\vec{e}_\beta}$. The duality between bases $\vec{e}^\nu \cdot \vec{e}_\alpha = \delta_\alpha^\nu$ then implies

$$\boxed{\nabla_\beta \vec{e}^\mu \equiv \nabla_{\vec{e}_\beta} \vec{e}^\mu = -\Gamma^\mu_{\alpha\beta} \vec{e}^\alpha}. \quad (23.31)$$

Note the sign flip, which is required to keep $\nabla_\beta(\vec{e}^\mu \cdot \vec{e}_\alpha) = 0$, and note that the differentiation index always goes last. Duality also implies that Eqs. (23.30) and (23.31) can be rewritten as

$$\Gamma^\mu_{\alpha\beta} = \vec{e}^\mu \cdot \nabla_\beta \vec{e}_\alpha = -\vec{e}_\alpha \cdot \nabla_\beta \vec{e}^\mu. \quad (23.32)$$

With the aid of these connection coefficients, we can evaluate the components $A_{\alpha;\beta}$ of the gradient of a vector field in any basis. We just compute

$$\begin{aligned} A^\mu_{;\beta} \vec{e}_\mu &= \nabla_\beta \vec{A} = \nabla_\beta(\vec{A}^\mu \vec{e}_\mu) = (\nabla_\beta A^\mu) \vec{e}_\mu + \vec{A}^\mu \nabla_\beta \vec{e}_\mu \\ &= A^\mu_{;\beta} \vec{e}_\mu + A^\mu \Gamma^\alpha_{\mu\beta} \vec{e}_\alpha \\ &= (A^\mu_{;\beta} + A^\alpha \Gamma^\mu_{\alpha\beta}) \vec{e}_\mu. \end{aligned} \quad (23.33)$$

In going from the first line to the second, we have used the notation

$$A^\mu_{;\beta} \equiv \partial_{\vec{e}_\beta} A^\mu; \quad (23.34)$$

i.e. the comma denotes the result of letting a basis vector act as a differential operator on the component of the vector. In going from the second line of (23.33) to the third, we have renamed some summed-over indices. By comparing the first and last expressions in Eq. (23.33), we conclude that

$$\boxed{A^\mu{}_{;\beta} = A^\mu{}_{,\beta} + A^\alpha \Gamma^\mu{}_{\alpha\beta}}. \quad (23.35)$$

The first term in this equation describes the changes in \vec{A} associated with changes of its component A^μ ; the second term *corrects for* artificial changes of A^μ that are induced by turning and length changes of the basis vector \vec{e}_μ . We shall use the short-hand terminology that the second term “corrects” the index μ

By a similar computation, we conclude that in any basis the covariant components of the gradient are

$$\boxed{A_{\alpha;\beta} = A_{\alpha,\beta} - \Gamma^\mu{}_{\alpha\beta} A_\mu}, \quad (23.36)$$

where again $A_{\alpha,\beta} \equiv \partial_{\vec{e}_\beta} A_\alpha$. Notice that when the index being “corrected” is down [α in Eq. (23.36)], the connection coefficient has a minus sign; when it is up [μ in Eq. (23.35)], the connection coefficient has a plus sign. This is in accord with the signs in Eqs. (23.31)–(23.32).

These considerations should make obvious the following equations for the components of the gradient of a tensor:

$$F^{\alpha\beta}{}_{;\gamma} = F^{\alpha\beta}{}_{,\gamma} + \Gamma^\alpha{}_{\mu\gamma} F^{\mu\beta} + \Gamma^\beta{}_{\mu\gamma} F^{\alpha\mu}, \quad F_{\alpha\beta;\gamma} = F_{\alpha\beta,\gamma} - \Gamma^\mu{}_{\alpha\gamma} F_{\mu\beta} - \Gamma^\mu{}_{\beta\gamma} F_{\alpha\mu}. \quad (23.37)$$

Notice that each index of \mathbf{F} must be corrected, the correction has a sign dictated by whether the index is up or down, the differentiation index always goes last on the Γ , and all other indices can be deduced by requiring that the free indices in each term be the same and all other indices be summed.

If we have been given a basis, then how can we compute the connection coefficients? We can try to do so by drawing pictures and examining how the basis vectors change from point to point—a method that is fruitful in spherical and cylindrical coordinates in Euclidean space (Sec. 11.5). However, in other situations this method is fraught with peril, so we need a firm mathematical prescription. It turns out that the following prescription works; see below for a proof:

(i) Evaluate the *commutation coefficients* $c_{\alpha\beta}{}^\rho$ of the basis, which are defined by the two equivalent relations

$$\boxed{[\vec{e}_\alpha, \vec{e}_\beta] \equiv c_{\alpha\beta}{}^\rho \vec{e}_\rho, \quad c_{\alpha\beta}{}^\rho \equiv \vec{e}^\rho \cdot [\vec{e}_\alpha, \vec{e}_\beta]}. \quad (23.38a)$$

[Note that in a coordinate basis the commutation coefficients will vanish. *Warning:* commutation coefficients also appear in the theory of Lie Groups; there it is conventional to use a different ordering of indices than here, $c_{\alpha\beta}{}^\rho{}_{\text{here}} = c^\rho{}_{\alpha\beta\text{Lie groups}}$.] (ii) Lower the last index on the commutation coefficients using the metric components in the basis:

$$\boxed{c_{\alpha\beta\gamma} \equiv c_{\alpha\beta}{}^\rho g_{\rho\gamma}}. \quad (23.38b)$$

(iii) Compute the quantities

$$\boxed{\Gamma_{\alpha\beta\gamma} \equiv \frac{1}{2}(g_{\alpha\beta,\gamma} + g_{\alpha\gamma,\beta} - g_{\beta\gamma,\alpha} + c_{\alpha\beta\gamma} + c_{\alpha\gamma\beta} - c_{\beta\gamma\alpha})}. \quad (23.38c)$$

Here the commas denote differentiation with respect to the basis vectors as though the metric components were scalar fields [as in Eq. (23.34)]. Notice that the pattern of indices is the same on the g 's and on the c 's. It is a peculiar pattern—one of the few aspects of index gymnastics that cannot be reconstructed by merely lining up indices. In a coordinate basis the c terms will vanish, so $\Gamma_{\alpha\beta\gamma}$ will be symmetric in its last two indices. In an orthonormal basis $g_{\mu\nu}$ are constant so the g terms will vanish, and $\Gamma_{\alpha\beta\gamma}$ will be antisymmetric in its first two indices. And in a Cartesian or Lorentz coordinate basis, which is both coordinate and orthonormal, both the c terms and the g terms will vanish, so $\Gamma_{\alpha\beta\gamma}$ will vanish. (iv) Raise the first index on $\Gamma_{\alpha\beta\gamma}$ to obtain the *connection coefficients*

$$\boxed{\Gamma^\mu{}_{\beta\gamma} = g^{\mu\alpha}\Gamma_{\alpha\beta\gamma}}. \quad (23.38d)$$

In a coordinate basis, the $\Gamma^\mu{}_{\beta\gamma}$ are called *Christoffel symbols*.

The above prescription, steps (i), (ii), (iii), for computing the connection coefficients follows from two key properties of the gradient $\vec{\nabla}$: *First*, The gradient of the metric tensor vanishes,

$$\boxed{\vec{\nabla}\mathbf{g} = 0}. \quad (23.39)$$

Second, for any two vector fields \vec{A} and \vec{B} , the gradient is related to the commutator by

$$\boxed{\nabla_{\vec{A}}\vec{B} - \nabla_{\vec{B}}\vec{A} = [\vec{A}, \vec{B}]}. \quad (23.40)$$

For a derivation of these relations and then a derivation of the prescription (i), (ii), (iii), see Exs. 23.6 and 23.7.

The gradient operator $\vec{\nabla}$ is an example of a geometric object that is not a tensor. The connection coefficients $\Gamma^\mu{}_{\beta\gamma} = \vec{e}^\mu \cdot (\nabla_{\vec{e}_\beta}\vec{e}_\gamma)$ can be regarded as the components of $\vec{\nabla}$; and because it is not a tensor, these components do not obey the tensorial transformation law (23.19) when switching from one basis to another. Their transformation law is far more complicated and is very rarely used. Normally one computes them from scratch in the new basis, using the above prescription or some other, equivalent prescription (cf. Chap. 14 of MTW). For most curved spacetimes that one meets in general relativity, these computations are long and tedious and therefore are normally carried out on computers using symbolic-manipulation software such as Maple or Mathematica, or programs such as GR-Tensor and MathTensor that run under Maple or Mathematica. Such software is easily found on the Internet using a search engine. A particularly simple Mathematica program for use with coordinate bases is presented and discussed in Appendix C of Hartle (2003), and is available on that book's website, <http://web.physics.ucsb.edu/~gravitybook/>.

EXERCISES

Exercise 23.6 *Derivation: Properties of the Gradient $\vec{\nabla}$*

- (a) Derive Eq. (23.39). [Hint: At a point \mathcal{P} where $\vec{\nabla}\mathbf{g}$ is to be evaluated, introduce a locally orthonormal coordinate basis (i.e. locally Cartesian or locally Lorentz). When computing in this basis, the effects of curvature will show up only at second order in distance from \mathcal{P} . Show that in this basis, the components of $\vec{\nabla}\mathbf{g}$ vanish, and from this infer that $\vec{\nabla}\mathbf{g}$, viewed as a frame-independent third-rank tensor, vanishes.]
- (b) Derive Eq. (23.40). [Hint: Again work in a locally orthonormal coordinate basis.]

Exercise 23.7 *Derivation: Prescription for Computing Connection Coefficients*

Derive the prescription (i), (ii), (iii) [Eqs. (23.38)] for computing the connection coefficients in any basis. [Hints: (i) In the chosen basis, from $\vec{\nabla}\mathbf{g} = 0$ infer that $\Gamma_{\alpha\beta\gamma} + \Gamma_{\beta\alpha\gamma} = g_{\alpha\beta,\gamma}$. Notice that this determines the part of $\Gamma_{\alpha\beta\gamma}$ that is symmetric in its first two indices. Show that the number of independent components of $\Gamma_{\alpha\beta\gamma}$ thereby determined is $\frac{1}{2}n^2(n+1)$. (ii) From Eq. (23.40) infer that $\Gamma_{\gamma\beta\alpha} - \Gamma_{\gamma\alpha\beta} = c_{\alpha\beta\gamma}$, which fixes the part of Γ antisymmetric in the last two indices. Show that the number of independent components thereby determined is $\frac{1}{2}n^2(n-1)$. (iii) Infer that the number of independent components determined by (i) and (ii) together is n^3 , which is the entirety of $\Gamma_{\alpha\beta\gamma}$. By somewhat complicated algebra, deduce Eq. (23.38c) for $\Gamma_{\alpha\beta\gamma}$. (The algebra is sketched in Ex. 8.15 of MTW). (iv) Then infer the final answer (23.38d) for $\Gamma^\mu{}_{\beta\gamma}$.]

Exercise 23.8 *Practice: Commutation and Connection Coefficients for Circular Polar Bases*

Consider the circular coordinates $\{\varpi, \phi\}$ of Fig. 23.3 and their associated bases.

- (a) Evaluate the commutation coefficients $c_{\alpha\beta}{}^\rho$ for the coordinate basis $\{\vec{e}_\varpi, \vec{e}_\phi\}$, and also for the orthonormal basis $\{\vec{e}_{\hat{\varpi}}, \vec{e}_{\hat{\phi}}\}$.
- (b) Compute by hand the connection coefficients for the coordinate basis and also for the orthonormal basis, using Eqs. (23.38a)–(23.38d). [Note: the answer for the orthonormal basis was worked out by a different method in our study of elasticity theory; Eq. (11.40).]
- (c) Repeat this computation using symbolic manipulation software on a computer.

Exercise 23.9 *Practice: Connection Coefficients for Spherical Polar Coordinates*

- (a) Consider spherical polar coordinates in 3-dimensional space and verify that the non-zero connection coefficients, assuming an orthonormal basis, are given by Eq. (11.41).
- (b) Repeat the exercise assuming a coordinate basis with

$$\mathbf{e}_r \equiv \frac{\partial}{\partial r}, \quad \mathbf{e}_\theta \equiv \frac{\partial}{\partial \theta}, \quad \mathbf{e}_\phi \equiv \frac{\partial}{\partial \phi}. \quad (23.41)$$

- (c) Repeat both computations using symbolic manipulation software on a computer.

Exercise 23.10 Practice: Index Gymnastics — Geometric Optics

This exercise gives the reader practice in formal manipulations that involve the gradient operator. In the geometric optics approximation (Chap. 7), for electromagnetic waves in Lorenz gauge one can write the 4-vector potential in the form $\vec{A} = \vec{\mathcal{A}}e^{i\varphi}$, where $\vec{\mathcal{A}}$ is a slowly varying amplitude and φ is a rapidly varying phase. By the techniques of Chap. 7, one can deduce from the vacuum Maxwell equations that the wave vector, defined by $\vec{k} \equiv \vec{\nabla}\varphi$, is null: $\vec{k} \cdot \vec{k} = 0$.

- (a) Rewrite all of the equations in the above paragraph in slot-naming index notation.
- (b) Using index manipulations, show that the wave vector \vec{k} (which is a vector field because the wave's phase φ is a scalar field) satisfies the geodesic equation, $\nabla_{\vec{k}}\vec{k} = 0$. The geodesics, to which \vec{k} is the tangent vector, are the rays discussed in Chap. 7, along which the waves propagate.

23.3.4 Integration

Our desire to use general bases and work in curved manifolds gives rise to two new issues in the definition of integrals.

First, the volume elements used in integration involve the Levi-Civita tensor [Eqs. (2.43), (2.52), (2.55)], so we need to know the components of the Levi-Civita tensor in a general basis. It turns out [see, e.g., Ex. 8.3 of MTW] that the covariant components differ from those in an orthonormal basis by a factor $\sqrt{|g|}$ and the contravariant by $1/\sqrt{|g|}$, where

$$\boxed{g \equiv \det ||g_{\alpha\beta}||} \quad (23.42)$$

is the determinant of the matrix whose entries are the covariant components of the metric. More specifically, let us denote by $[\alpha\beta\dots\nu]$ the value of $\epsilon_{\alpha\beta\dots\nu}$ in an orthonormal basis of our n -dimensional space [Eq. (2.43b)]:

$$\begin{aligned} [12\dots n] &= +1, \\ [\alpha\beta\dots\nu] &= +1 \text{ if } \alpha, \beta, \dots, \nu \text{ is an even permutation of } 1, 2, \dots, n \\ &= -1 \text{ if } \alpha, \beta, \dots, \nu \text{ is an odd permutation of } 1, 2, \dots, n \\ &= 0 \text{ if } \alpha, \beta, \dots, \nu \text{ are not all different.} \end{aligned} \quad (23.43)$$

(In spacetime the indices must run from 0 to 3 rather than 1 to $n = 4$). Then in a general right-handed basis the components of the Levi-Civita tensor are

$$\boxed{\epsilon_{\alpha\beta\dots\nu} = \sqrt{|g|} [\alpha\beta\dots\nu], \quad \epsilon^{\alpha\beta\dots\nu} = \pm \frac{1}{\sqrt{|g|}} [\alpha\beta\dots\nu]}, \quad (23.44)$$

where the \pm is plus in Euclidean space and minus in spacetime. In a left-handed basis the sign is reversed.

As an example of these formulas, consider a spherical polar coordinate system (r, θ, ϕ) in three-dimensional Euclidean space, and use the three infinitesimal vectors $dx^j(\partial/\partial x^j)$ to construct the volume element $d\Sigma$ [cf. Eq. (1.27)]:

$$d\Sigma = \epsilon \left(dr \frac{\partial}{\partial r}, d\theta \frac{\partial}{\partial \theta}, d\phi \frac{\partial}{\partial \phi} \right) = \epsilon_{r\theta\phi} dr d\theta d\phi = \sqrt{g} dr d\theta d\phi = r^2 \sin \theta dr d\theta d\phi. \quad (23.45)$$

Here the second equality follows from linearity of ϵ and the formula for computing its components by inserting basis vectors into its slots; the third equality follows from our formula (23.44) for the components, and the fourth equality entails the determinant of the metric coefficients, which in spherical coordinates are $g_{rr} = 1$, $g_{\theta\theta} = r^2$, $g_{\phi\phi} = r^2 \sin^2 \theta$, all other g_{jk} vanish, so $g = r^4 \sin^2 \theta$. The resulting volume element $r^2 \sin \theta dr d\theta d\phi$ should be familiar and obvious.

The *second* new integration issue that we must face is the fact that integrals such as

$$\int_{\partial\mathcal{V}} T^{\alpha\beta} d\Sigma_\beta \quad (23.46)$$

[cf. Eqs. (2.55), (2.56)] involve constructing a vector $T^{\alpha\beta} d\Sigma_\beta$ in each infinitesimal region $d\Sigma_\beta$ of the surface of integration, and then adding up the contributions from all the infinitesimal regions. A major difficulty arises from the fact that each contribution lives in a different tangent space. To add them together, we must first transport them all to the same tangent space at some single location in the manifold. How is that transport to be performed? The obvious answer is “by the same parallel transport technique that we used in defining the gradient.” However, when defining the gradient we only needed to perform the parallel transport over an infinitesimal distance, and now we must perform it over long distances. As we shall see in Chap. 24, when the manifold is curved, long-distance parallel transport gives a result that depends on the route of the transport, and in general there is no way to identify any preferred route.

As a result, *integrals such as (23.46) are ill-defined in a curved manifold. The only integrals that are well defined in a curved manifold are those such as $\int_{\partial\mathcal{V}} S^\alpha d\Sigma_\alpha$ whose infinitesimal contributions $S^\alpha d\Sigma_\alpha$ are scalars, i.e. integrals whose value is a scalar.* This fact will have profound consequences in curved spacetime for the laws of conservation of energy, momentum, and angular momentum (Secs. 24.7 and 24.9.4).

EXERCISES

Exercise 23.11 *Practice: Integration — Gauss’s Theorem*

In 3-dimensional Euclidean space the Maxwell equation $\nabla \cdot \mathbf{E} = \rho_e/\epsilon_0$ can be combined with Gauss’s theorem to show that the electric flux through the surface $\partial\mathcal{V}$ of a sphere is equal to the charge in the sphere’s interior \mathcal{V} divided by ϵ_0 :

$$\int_{\partial\mathcal{V}} \mathbf{E} \cdot d\Sigma = \int_{\mathcal{V}} (\rho_e/\epsilon_0) d\Sigma. \quad (23.47)$$

Introduce spherical polar coordinates so the sphere's surface is at some radius $r = R$. Consider a surface element on the sphere's surface with vectorial legs $d\phi\partial/\partial\phi$ and $d\theta\partial/\partial\theta$. Evaluate the components $d\Sigma_j$ of the surface integration element $d\Sigma = \epsilon(\dots, d\theta\partial/\partial\theta, d\phi\partial/\partial\phi)$. (Here ϵ is the Levi-Civita tensor.) Similarly, evaluate $d\Sigma$ in terms of vectorial legs in the sphere's interior. Then use these results for $d\Sigma_j$ and $d\Sigma$ to convert Eq. (23.47) into an explicit form in terms of integrals over r, θ, ϕ . The final answer should be obvious, but the above steps in deriving it are informative.

23.4 The Stress-Energy Tensor Revisited

In Sec. 2.13.1, we defined the *stress-energy tensor* \mathbf{T} of any matter or field as a symmetric, second-rank tensor that describes the flow of 4-momentum through spacetime. More specifically, the total 4-momentum \vec{P} that flows through some small 3-volume $\vec{\Sigma}$ (defined in Sec. 2.12.1), going from the negative side of $\vec{\Sigma}$ to its positive side, is

$$\mathbf{T}(_, \vec{\Sigma}) = (\text{total 4-momentum } \vec{P} \text{ that flows through } \vec{\Sigma}); \quad \text{i.e., } T^{\alpha\beta}\Sigma_\beta = P^\alpha \quad (23.48)$$

[Eq. (2.66)]. Of course, this stress-energy tensor depends on the location \mathcal{P} of the 3-volume in spacetime; i.e., it is a tensor field $\mathbf{T}(\mathcal{P})$.

From this geometric, frame-independent definition of the stress-energy tensor, we were able to read off the physical meaning of its components in any inertial reference frame [Eqs. (2.67)]: T^{00} is the total energy density, including rest mass-energy; $T^{j0} = T^{0j}$ is the j -component of momentum density, or equivalently the j -component of energy flux; and T^{jk} are the components of the stress tensor, or equivalently of the momentum flux.

In Sec. 2.13.2, we formulated the law of conservation of 4-momentum in a global form and a local form. The global form, $\int_{\partial\mathcal{V}} T^{\alpha\beta} d\Sigma_\beta = 0$ [Eq. (2.71)] says that all the 4-momentum that enters a closed 4-volume \mathcal{V} in spacetime through its boundary $\partial\mathcal{V}$ in the past must ultimately exit through $\partial\mathcal{V}$ in the future; cf. Fig. 2.12. The local form of 4-momentum conservation,

$$\vec{\nabla} \cdot \mathbf{T} = 0, \quad (23.49)$$

says that, in any chosen Lorentz frame, the time derivative of the energy density plus the divergence of the energy flux vanishes, $\partial T^{00}/\partial t + \partial T^{0j}/\partial x^j = 0$ and similarly for the momentum, $\partial T^{j0}/\partial t + \partial T^{jk}/\partial x^k = 0$.

The stress-energy tensor and 4-momentum conservation will play a major role in our development of general relativity. Almost all of our examples will entail perfect fluids.

Recall [Eq. (2.74a)] that in the local rest frame of a perfect fluid, there is no energy flux or momentum density, $T^{j0} = T^{0j} = 0$, but there is a total energy density (including rest mass) ρ and an isotropic pressure P :

$$T^{00} = \rho, \quad T^{jk} = P\delta^{jk}. \quad (23.50)$$

From this special form of $T^{\alpha\beta}$ in the fluid's local rest frame, one can derive a geometric, frame-independent expression for the fluid's stress-energy tensor \mathbf{T} in terms of its 4-velocity \vec{u} , the metric tensor \mathbf{g} , and the rest-frame energy density ρ and pressure P :

$$\boxed{\mathbf{T} = (\rho + P)\vec{u} \otimes \vec{u} + P\mathbf{g}; \quad \text{i.e., } T^{\alpha\beta} = (\rho + P)u^\alpha u^\beta + P g^{\alpha\beta}}; \quad (23.51)$$

[Eq. (2.74b)]; see Ex. 2.26. This expression for the stress-energy tensor of a perfect fluid is an example of a geometric, frame-independent description of physics.

The equations of relativistic fluid dynamics for a perfect fluid are obtained by inserting the stress-energy tensor (23.51) into the law of 4-momentum conservation $\vec{\nabla} \cdot \mathbf{T} = 0$, and augmenting with the law of rest-mass conservation. We explored this in brief in Ex. 2.26, and in much greater detail in Sec. 13.8.2. Applications that we have explored are the relativistic Bernoulli equation and ultrarelativistic jets (Sec. 13.8.2) and relativistic shocks (Ex. 17.11). In Sec. 13.8.3 we explored in detail the slightly subtle way in which a fluid's nonrelativistic energy density, energy flux, and stress tensor arise from the relativistic perfect-fluid stress-energy tensor (23.51).

These issues for a perfect fluid are so important that readers are encouraged to review them (except possibly the applications) in preparation for our foray into general relativity.

Four other important examples of the stress-energy tensor are those for the electromagnetic field (Ex. 2.28), for a kinetic-theory swarm of relativistic particles (Secs. 3.4.2 and 3.5.3), for a point particle (Box 23.2) and for a relativistic fluid with viscosity and diffusive heat conduction (Ex. 23.13). However, we shall not do much with any of these during our study of general relativity.

EXERCISES

Exercise 23.12 $T2$ *Derivation: Stress-Energy Tensor for a Point Particle*
Derive Eq. (3) of Box 23.2.

Exercise 23.13 *Example: Stress-Energy Tensor for a Viscous Fluid with Diffusive Heat Conduction*

This exercise serves two roles: It develops the relativistic stress-energy tensor for a viscous fluid with diffusive heat conduction, and in the process it allows the reader to gain practice in index gymnastics.

In our study of elasticity theory, we introduced the concept of the irreducible tensorial parts of a second-rank tensor in Euclidean space (Box. 11.2). Consider a relativistic fluid flowing through spacetime, with a 4-velocity $\vec{u}(\mathcal{P})$. The fluid's gradient $\vec{\nabla}\vec{u}$ ($u_{\alpha;\beta}$ in slot-naming index notation) is a second-rank tensor in spacetime. With the aid of the 4-velocity itself, we can break it down into irreducible tensorial parts as follows:

$$u_{\alpha;\beta} = -a_\alpha u_\beta + \frac{1}{3}\theta P_{\alpha\beta} + \sigma_{\alpha\beta} + \omega_{\alpha\beta}. \quad (23.52)$$

Box 23.2

T2 Stress-Energy Tensor for a Point Particle

For a point particle which moves through spacetime along a world line $\mathcal{P}(\zeta)$ (where ζ is the affine parameter such that the particle's 4-momentum is $\vec{p} = d/d\zeta$), the stress-energy tensor will vanish everywhere except on the world line itself. Correspondingly, \mathbf{T} must be expressed in terms of a Dirac delta function. The relevant delta function is a scalar function of two points in spacetime, $\delta(\mathcal{Q}, \mathcal{P})$ with the property that when one integrates over the point \mathcal{P} , using the 4-dimensional volume element $d\Sigma$ (which in any inertial frame just reduces to $d\Sigma = dt dx dy dz$), one obtains

$$\int_{\mathcal{V}} f(\mathcal{P}) \delta(\mathcal{Q}, \mathcal{P}) d\Sigma = f(\mathcal{Q}) . \quad (1)$$

Here $f(\mathcal{P})$ is an arbitrary scalar field and the region \mathcal{V} of 4-dimensional integration must include the point \mathcal{Q} . One can easily verify that in terms of Lorentz coordinates this delta function can be expressed as

$$\delta(\mathcal{Q}, \mathcal{P}) = \delta(t_{\mathcal{Q}} - t_{\mathcal{P}}) \delta(x_{\mathcal{Q}} - x_{\mathcal{P}}) \delta(y_{\mathcal{Q}} - y_{\mathcal{P}}) \delta(z_{\mathcal{Q}} - z_{\mathcal{P}}) , \quad (2)$$

where the deltas on the right-hand side are ordinary one-dimensional Dirac delta functions. [Proof: Simply insert Eq. (2) into Eq. (1), replace $d\Sigma$ by $dt_{\mathcal{Q}} dx_{\mathcal{Q}} dy_{\mathcal{Q}} dz_{\mathcal{Q}}$, and perform the four integrations.]

The general definition (23.48) of the stress-energy tensor \mathbf{T} implies that that the integral of a point particle's stress-energy tensor over any 3-surface \mathcal{S} that slices through the particle's world line just once, at an event $\mathcal{P}(\zeta_o)$, must be equal to the particle's 4-momentum at the intersection point:

$$\int_{\mathcal{S}} T^{\alpha\beta} d\Sigma_{\beta} = p^{\alpha}(\zeta_o) . \quad (3)$$

It is a straightforward but sophisticate exercise [Ex. 23.12] to verify that the following frame-independent expression has this property:

$$\mathbf{T}(\mathcal{Q}) = \int_{-\infty}^{+\infty} \vec{p}(\zeta) \otimes \vec{p}(\zeta) \delta(\mathcal{Q}, \mathcal{P}(\zeta)) d\zeta . \quad (4)$$

Here the integral is along the world line $\mathcal{P}(\zeta)$ of the particle, and \mathcal{Q} is the point at which \mathbf{T} is being evaluated [the integration point in Eq. (3)]. Therefore, Eq. (4) is the point-particle stress-energy tensor.

Here: (i)

$$P_{\alpha\beta} \equiv g_{\alpha\beta} + u_\alpha u_\beta \quad (23.53)$$

is a tensor that projects vectors into the 3-space orthogonal to \vec{u} ; it can also be regarded as that 3-space's metric; see Ex. 2.10. (ii) $\sigma_{\alpha\beta}$ is symmetric and trace-free and is orthogonal to the 4-velocity, and (iii) $\omega_{\alpha\beta}$ is antisymmetric and is orthogonal to the 4-velocity.

- (a) Show that the rate of change of \vec{u} along itself, $\nabla_{\vec{u}}\vec{u}$ (i.e., the fluid 4-acceleration) is equal to the vector \vec{a} that appears in the decomposition (23.52). Show, further, that $\vec{a} \cdot \vec{u} = 0$.
- (b) Show that the divergence of the 4-velocity, $\nabla \cdot \vec{u}$, is equal to the scalar field θ that appears in the decomposition (23.52).
- (c) The quantities $\sigma_{\alpha\beta}$ and $\omega_{\alpha\beta}$ are the relativistic versions of a Newtonian fluid's shear and rotation tensors, which we introduced in Sec. 13.7.1. Derive equations for these tensors in terms of $u_{\alpha;\beta}$ and $P_{\mu\nu}$.
- (d) Show that, as viewed in a Lorentz reference frame where the fluid is moving with speed small compared to the speed of light, to first-order in the fluid's ordinary velocity $v^j = dx^j/dt$, the following are true: (i) $u^0 = 1$, $u^j = v^j$; (ii) θ is the nonrelativistic expansion of the fluid, $\theta = \nabla \cdot \mathbf{v} \equiv v^j_{,j}$ [Eq. (13.63)]; (iii) σ_{jk} is the fluid's nonrelativistic shear [Eq. (13.63)]; (iv) ω_{jk} is the fluid's nonrelativistic rotation tensor [denoted r_{jk} in Eq. (13.63)].
- (e) At some event \mathcal{P} where we want to know the influence of viscosity on the fluid's stress-energy tensor, introduce the fluid's local rest frame. Explain why, in that frame, the only contributions of viscosity to the components of the stress-energy tensor are $T_{\text{visc}}^{jk} = -\zeta\theta g^{jk} - 2\mu\sigma^{jk}$, where ζ and μ are the coefficient's of bulk and shear viscosity; the contributions to T^{00} and $T^{j0} = T^{0j}$ vanish. [Hint: see Eq. (13.68) and associated discussions].
- (f) From nonrelativistic fluid mechanics, infer that, in the fluid's rest frame at \mathcal{P} , the only contributions of diffusive heat conductivity to the stress-energy tensor are $T_{\text{cond}}^{0j} = T_{\text{cond}}^{j0} = -\kappa\partial T/\partial x^j$, where κ is the fluid's thermal conductivity and T is its temperature. [Hint: see Eq. (13.69) and associated discussion.]
- (g) From this, deduce the following geometric, frame-invariant form of the fluid's stress-energy tensor:

$$T_{\alpha\beta} = (\rho + P)u_\alpha u_\beta + P g_{\alpha\beta} - \zeta\theta g_{\alpha\beta} - 2\mu\sigma_{\alpha\beta} - 2\kappa u_{(\alpha} P_{\beta)}{}^{\mu} T_{;\mu} . \quad (23.54)$$

Here the parentheses in the last term mean to symmetrize in the α and β slots.

Comment: From the divergence of this stress-energy tensor, plus the first law of thermodynamics and the law of rest-mass conservation, one can derive the full theory of relativistic fluid mechanics for a fluid with viscosity and heat flow; see, e.g., Ex. 22.7 of MTW.

23.5 The Proper Reference Frame of an Accelerated Observer

Physics experiments and astronomical measurements almost always use apparatus that accelerates and rotates. For example, if the apparatus is in an earth-bound laboratory and is attached to the laboratory floor and walls, then it accelerates upward (relative to freely falling particles) with the negative of the “acceleration of gravity”, and it rotates (relative to inertial gyroscopes) because of the rotation of the earth. It is useful, in studying such apparatus, to regard it as attached to an accelerating, rotating reference frame. As preparation for studying such reference frames in the presence of gravity, we here shall study them in flat spacetime. For a somewhat more sophisticated treatment, see pages 163–176, 327–332 of MTW.

Consider an observer with 4-velocity \vec{U} , who moves along an accelerated world line through flat spacetime (Fig. 23.6) so she has a nonzero 4-acceleration

$$\vec{a} = \vec{\nabla}_{\vec{U}} \vec{U} . \quad (23.55)$$

Have that observer construct, in the vicinity of her world line, a coordinate system $\{x^{\hat{\alpha}}\}$ (called her *proper reference frame*) with these properties: (i) The spatial origin is centered on her world line at all times, i.e., her world line is given by $x^{\hat{j}} = 0$. (ii) Along her world line, the time coordinate $x^{\hat{0}}$ is the same as the proper time ticked by an ideal clock that she carries. (iii) In the immediate vicinity of her world line, the spatial coordinates $x^{\hat{j}}$ measure




Fig. 23.6: The proper reference frame of an accelerated observer. The spatial basis vectors $\vec{e}_{\hat{x}}$, $\vec{e}_{\hat{y}}$, and $\vec{e}_{\hat{z}}$ are orthogonal to the observer’s world line and rotate, relative to local gyroscopes, as they move along the world line. The flat 3-planes spanned by these basis vectors are surfaces of constant coordinate time $x^{\hat{0}} \equiv$ (proper time as measured by the observer’s clock at the event where the 3-plane intersects the observer’s world line); in other words, they are the observer’s “3-space”. In each of these flat 3-planes the spatial coordinates \hat{x} , \hat{y} , \hat{z} are Cartesian, with $\partial/\partial\hat{x} = \vec{e}_{\hat{x}}$, $\partial/\partial\hat{y} = \vec{e}_{\hat{y}}$, $\partial/\partial\hat{z} = \vec{e}_{\hat{z}}$.

physical distance along the axes of a little Cartesian latticework that she carries (and that she regards as purely spatial, which means it lies in the 3-plane orthogonal to her world line). These properties dictate that, in the immediate vicinity of her world line, the metric has the form $ds^2 = \eta_{\hat{\alpha}\hat{\beta}} dx^{\hat{\alpha}} dx^{\hat{\beta}}$, where $\eta_{\hat{\alpha}\hat{\beta}}$ are the Lorentz-basis metric coefficients, Eq. (23.6); in other words, all along her world line the coordinate basis vectors are orthonormal:

$$g_{\hat{\alpha}\hat{\beta}} = \frac{\partial}{\partial x^{\hat{\alpha}}} \cdot \frac{\partial}{\partial x^{\hat{\beta}}} = \eta_{\hat{\alpha}\hat{\beta}} \quad \text{at } x^{\hat{j}} = 0. \quad (23.56)$$

Properties (i) and (ii) dictate, moreover, that along the observer's world line, the basis vector $\vec{e}_{\hat{0}} \equiv \partial/\partial x^{\hat{0}}$ differentiates with respect to her proper time, and thus is identically equal to her 4-velocity \vec{U} ,

$$\vec{e}_{\hat{0}} = \frac{\partial}{\partial x^{\hat{0}}} = \vec{U}. \quad (23.57)$$

There remains freedom as to how the observer's latticework is oriented, spatially. The observer can lock it to the gyroscopes of an *inertial-guidance system* that she carries (Box 23.3), in which case we shall say that it is “nonrotating”; or she can rotate it relative to such gyroscopes. For generality, we shall assume that the latticework rotates. Its angular velocity, as measured by the observer (by comparing the latticework's orientation with inertial-guidance gyroscopes), is a 3-dimensional, spatial vector $\boldsymbol{\Omega}$ in the 3-plane orthogonal to her world line; and as viewed in 4-dimensional spacetime, it is a 4-vector $\vec{\Omega}$ whose components in the observer's reference frame are $\Omega^{\hat{j}} \neq 0$ and $\Omega^{\hat{0}} = 0$. Similarly, the latticework's acceleration, as measured by an inertial-guidance accelerometer attached to it (Box 23.3), is a 3-dimensional spatial vector \mathbf{a} which can be thought of as a 4-vector with components in

Box 23.3

Inertial Guidance Systems

Aircraft and rockets often carry inertial guidance systems, which consist of an accelerometer and a set of gyroscopes.

The accelerometer measures the system's 4-acceleration \vec{a} (in relativistic language). Equivalently, it measures the system's Newtonian 3-acceleration \mathbf{a} relative to inertial coordinates in which the system is momentarily at rest. As we shall see in Eq. (23.58) below, these quantities are two different ways of thinking about the same thing.

Each gyroscope is forced to remain at rest in the aircraft or rocket by a force that is applied at its center of mass. Such a force exerts no torque around the center of mass, so the gyroscope maintains its direction (does not precess) relative to an inertial frame in which it is momentarily at rest.

As the accelerating aircraft or rocket turns, its walls rotate with some angular velocity $\vec{\Omega}$ relative to these inertial-guidance gyroscopes. This is the angular velocity discussed in the text, between Eq. (23.57) and Eq. (23.58).

From the time evolving 4-acceleration $\vec{a}(\tau)$ and angular velocity $\vec{\Omega}(\tau)$, a computer can calculate the aircraft's (or rocket's) world line and its changing orientation.

the observer's frame

$$a^{\hat{0}} = 0, \quad a^{\hat{j}} = (\hat{j}\text{-component of the measured } \mathbf{a}). \quad (23.58)$$

This 4-vector, in fact, is the observer's 4-acceleration, as one can verify by computing the 4-acceleration in an inertial frame in which the observer is momentarily at rest.

Geometrically, the coordinates of the proper reference frame are constructed as follows: (i) Begin with the basis vectors $\vec{e}_{\hat{\alpha}}$ along the observer's world line (Fig. 23.6)—basis vectors that satisfy equations (23.56) and (23.57), and that rotate with angular velocity $\vec{\Omega}$ relative to gyroscopes. Through the observer's world line at time $x^{\hat{0}}$ construct the flat 3-plane spanned by the spatial basis vectors $\vec{e}_{\hat{j}}$. Because $\vec{e}_{\hat{j}} \cdot \vec{e}_{\hat{0}} = 0$, this 3-plane is orthogonal to the world line. All events in this 3-plane are given the same value of coordinate time $x^{\hat{0}}$ as the event where it intersects the world line; thus the 3-plane is a surface of constant coordinate time $x^{\hat{0}}$. The spatial coordinates in this flat 3-plane are ordinary, Cartesian coordinates $x^{\hat{j}}$ with $\vec{e}_{\hat{j}} = \partial/\partial x^{\hat{j}}$.

23.5.1 Relation to Inertial Coordinates; Metric in Proper Reference Frame; Transport Law for Rotating Vectors

It is instructive to examine the coordinate transformation between these proper-reference-frame coordinates $x^{\hat{\alpha}}$ and the coordinates x^{μ} of an inertial reference frame. We shall pick a very special inertial frame for this purpose: Choose an event on the observer's world line, near which the coordinate transformation is to be constructed; adjust the origin of the observer's proper time so this event is $x^{\hat{0}} = 0$ (and of course $x^{\hat{j}} = 0$); and choose the inertial frame to be one which, arbitrarily near this event, coincides with the observer's proper reference frame. Then, if we were doing Newtonian physics, the coordinate transformation from the proper reference frame to the inertial frame would have the form (accurate through terms quadratic in $x^{\hat{\alpha}}$)

$$x^i = x^{\hat{i}} + \frac{1}{2}a^{\hat{i}}(x^{\hat{0}})^2 + \epsilon^{\hat{i}}_{\hat{j}\hat{k}}\Omega^{\hat{j}}x^{\hat{k}}x^{\hat{0}}, \quad x^0 = x^{\hat{0}}. \quad (23.59)$$

Here the term $\frac{1}{2}a^{\hat{i}}(x^{\hat{0}})^2$ is the standard expression for the vectorial displacement produced, after time $x^{\hat{0}}$ by the acceleration $a^{\hat{j}}$; and the term $\epsilon^{\hat{i}}_{\hat{j}\hat{k}}\Omega^{\hat{j}}x^{\hat{k}}x^{\hat{0}}$ is the standard expression for the displacement produced by the rotation $\Omega^{\hat{j}}$ during a short time $x^{\hat{0}}$. In relativity theory there is only one departure from these familiar expressions (up through quadratic order): after time $x^{\hat{0}}$ the acceleration has produced a velocity $v^{\hat{j}} = a^{\hat{j}}x^{\hat{0}}$ of the proper reference frame relative to the inertial frame; and correspondingly there is a Lorentz-boost correction to the transformation of time: $x^0 = x^{\hat{0}} + v^{\hat{j}}x^{\hat{j}} = x^{\hat{0}}(1 + a_{\hat{j}}x^{\hat{j}})$ [cf. Eq. (2.37c)], accurate only to quadratic order. Thus, the full transformation to quadratic order is

$$\begin{aligned} x^i &= x^{\hat{i}} + \frac{1}{2}a^{\hat{i}}(x^{\hat{0}})^2 + \epsilon^{\hat{i}}_{\hat{j}\hat{k}}\Omega^{\hat{j}}x^{\hat{k}}x^{\hat{0}}, \\ x^0 &= x^{\hat{0}}(1 + a_{\hat{j}}x^{\hat{j}}). \end{aligned} \quad (23.60a)$$

From this transformation and the form of the metric, $ds^2 = -(dx^0)^2 + \delta_{ij}dx^i dx^j$ in the inertial frame, we easily can evaluate the form of the metric, accurate to linear order in \mathbf{x} ,

in the proper reference frame:

$$\boxed{ds^2 = -(1 + 2\mathbf{a} \cdot \mathbf{x})(dx^{\hat{0}})^2 + 2(\boldsymbol{\Omega} \times \mathbf{x}) \cdot d\mathbf{x} dx^{\hat{0}} + \delta_{jk} dx^{\hat{j}} dx^{\hat{k}}} \quad (23.60b)$$

[Ex. 23.14a]. Here the notation is that of 3-dimensional vector analysis, with \mathbf{x} the 3-vector whose components are $x^{\hat{j}}$, $d\mathbf{x}$ that with components $dx^{\hat{j}}$, \mathbf{a} that with components $a^{\hat{j}}$, and $\boldsymbol{\Omega}$ that with components $\Omega^{\hat{j}}$.

Because the transformation (23.60a) was constructed near an arbitrary event on the observer's world line, the metric (23.60b) is valid near any and every event on its world line; i.e., it is valid all along the world line. It, in fact, is the leading order in an expansion in powers of the spatial separation $x^{\hat{j}}$ from the world line. For higher order terms in this expansion see, e.g., Ni and Zimmermann (1978).

Notice that precisely on the observer's world line, the metric coefficients $g_{\hat{\alpha}\hat{\beta}}$ [the coefficients of $dx^{\hat{\alpha}} dx^{\hat{\beta}}$ in Eq. (23.60b)] are $g_{\hat{\alpha}\hat{\beta}} = \eta_{\hat{\alpha}\hat{\beta}}$, in accord with equation (23.56). However, as one moves farther and farther away from the observer's world line, the effects of the acceleration $a^{\hat{j}}$ and rotation $\Omega^{\hat{j}}$ cause the metric coefficients to deviate more and more strongly from $\eta_{\hat{\alpha}\hat{\beta}}$.

From the metric coefficients of Eq. (23.60b), one can compute the connection coefficients $\Gamma^{\hat{\alpha}}_{\hat{\beta}\hat{\gamma}}$ on the observer's world line; and from these connection coefficients, one can infer the rates of change of the basis vectors along the world line, $\nabla_{\vec{U}} \vec{e}_{\hat{\alpha}} = \nabla_{\hat{0}} \vec{e}_{\hat{\alpha}} = \Gamma^{\hat{\mu}}_{\hat{\alpha}\hat{0}} \vec{e}_{\hat{\mu}}$. The result is [Ex. 23.14]

$$\nabla_{\vec{U}} \hat{e}_{\hat{0}} \equiv \nabla_{\vec{U}} \vec{U} = \vec{a}, \quad (23.61a)$$

$$\nabla_{\vec{U}} \vec{e}_{\hat{j}} = (\vec{a} \cdot \vec{e}_{\hat{j}}) \vec{U} + \boldsymbol{\epsilon}(\vec{U}, \vec{\Omega}, \vec{e}_{\hat{j}}, \underline{\quad}). \quad (23.61b)$$

Equation (23.61b) is general “law of transport” for constant-length vectors that are orthogonal to the observer's world line and that the observer thus sees as purely spatial: For the spin vector \vec{S} of an inertial-guidance gyroscope (Box 23.3), the transport law is (23.61b) with $\vec{e}_{\hat{j}}$ replaced by \vec{S} and with $\vec{\Omega} = 0$:

$$\boxed{\nabla_{\vec{U}} \vec{S} = \vec{U}(\vec{a} \cdot \vec{S})}; \quad (23.62)$$

This is called *Fermi-Walker transport*. The term on the right-hand side of this transport law is required to keep the spin vector always orthogonal to the observer's 4-velocity, $\nabla_{\vec{U}}(\vec{S} \cdot \vec{U}) = 0$. For any other vector \vec{A} , which rotates relative to inertial-guidance gyroscopes, the transport law has in addition to this “keep-it-orthogonal-to \vec{U} ” term, also a second term which is the 4-vector form of $d\mathbf{A}/dt = \boldsymbol{\Omega} \times \mathbf{A}$:

$$\nabla_{\vec{U}} \vec{A} = \vec{U}(\vec{a} \cdot \vec{A}) + \boldsymbol{\epsilon}(\vec{U}, \vec{\Omega}, \vec{A}, \dots). \quad (23.63)$$

Equation (23.61b) is this general transport law with \vec{A} replaced by $\vec{e}_{\hat{j}}$.

23.5.2 Geodesic Equation for Freely Falling Particle

Consider a particle with 4-velocity \vec{u} that moves freely through the neighborhood of an accelerated observer. As seen in an inertial reference frame, the particle moves through spacetime on a straight line, also called a *geodesic* of flat spacetime. Correspondingly, a geometric, frame-independent version of its *geodesic law of motion* is

$$\boxed{\nabla_{\vec{u}}\vec{u} = 0} ; \quad (23.64)$$

i.e., it parallel transports its 4-velocity \vec{u} along itself. It is instructive to examine the component form of this “geodesic equation” in the proper reference frame of the observer. Since the components of \vec{u} in this frame are $u^\alpha = dx^\alpha/d\tau$, where τ is the particle’s proper time (not the observer’s proper time), the components $u^{\hat{\alpha}}{}_{;\hat{\mu}}u^{\hat{\mu}} = 0$ of the geodesic equation (23.64) are

$$u^{\hat{\alpha}}{}_{;\hat{\mu}}u^{\hat{\mu}} + \Gamma^{\hat{\alpha}}{}_{\hat{\mu}\hat{\nu}}u^{\hat{\mu}}u^{\hat{\nu}} = \left(\frac{\partial}{\partial x^{\hat{\mu}}} \frac{dx^{\hat{\alpha}}}{d\tau} \right) \frac{dx^{\hat{\mu}}}{d\tau} + \Gamma^{\hat{\alpha}}{}_{\hat{\mu}\hat{\nu}}u^{\hat{\mu}}u^{\hat{\nu}} = 0 ; \quad (23.65)$$

or equivalently

$$\boxed{\frac{d^2x^{\hat{\alpha}}}{d\tau^2} + \Gamma^{\hat{\alpha}}{}_{\hat{\mu}\hat{\nu}} \frac{dx^{\hat{\mu}}}{d\tau} \frac{dx^{\hat{\nu}}}{d\tau} = 0} . \quad (23.66)$$

Suppose, for simplicity, that the particle is moving slowly relative to the observer, so its ordinary velocity $v^{\hat{j}} = dx^{\hat{j}}/dx^{\hat{0}}$ is very nearly equal to $u^{\hat{j}} = dx^{\hat{j}}/d\tau$ and is very small compared to unity (the speed of light), and $u^{\hat{0}} = dx^{\hat{0}}/d\tau$ is very nearly unity. Then to first order in the ordinary velocity $v^{\hat{j}}$, the spatial part of the geodesic equation (23.66) becomes

$$\frac{d^2x^{\hat{i}}}{(dx^{\hat{0}})^2} = -\Gamma^{\hat{i}}{}_{\hat{0}\hat{0}} - (\Gamma^{\hat{i}}{}_{\hat{j}\hat{0}} + \Gamma^{\hat{i}}{}_{\hat{0}\hat{j}})v^{\hat{j}} . \quad (23.67)$$

By computing the connection coefficients from the metric coefficients of Eq. (23.60b) [Ex. 23.14], we bring this low-velocity geodesic law of motion into the form

$$\frac{d^2x^{\hat{i}}}{(dx^{\hat{0}})^2} = -a^{\hat{i}} - 2\epsilon^{\hat{i}}{}_{\hat{j}\hat{k}}\Omega^{\hat{j}}v^{\hat{k}} , \quad \text{i.e.,} \quad \frac{d^2\mathbf{x}}{(dx^{\hat{0}})^2} = -\mathbf{a} - 2\boldsymbol{\Omega} \times \mathbf{v} . \quad (23.68)$$

This is the standard nonrelativistic form of the law of motion for a free particle as seen in a rotating, accelerating reference frame: the first term on the right-hand side is the inertial acceleration due to the failure of the frame to fall freely, and the second term is the Coriolis acceleration due to the frame’s rotation. There would also be a centrifugal acceleration if we had kept terms higher order in distance away from the observer’s world line, but it has been lost due to our linearizing the metric (23.60b) in that distance.

This analysis shows how the elegant formalism of tensor analysis gives rise to familiar physics. In the next few chapters we will see it give rise to less familiar, general relativistic phenomena.

EXERCISES

Exercise 23.14 *Derivation: Proper Reference Frame*

- (a) Show that the coordinate transformation (23.60a) brings the metric $ds^2 = \eta_{\alpha\beta} dx^\alpha dx^\beta$ into the form (23.60b), accurate to linear order in separation $x^{\hat{j}}$ from the origin of coordinates.
- (b) Compute the connection coefficients for the coordinate basis of (23.60b) at an arbitrary event on the observer's world line. Do so first by hand calculations, and then verify your results using symbolic-manipulation software on a computer.
- (c) From those connection coefficients show that the rate of change of the basis vectors $\mathbf{e}_{\hat{a}}$ along the observer's world line is given by (23.61a), (23.61b).
- (d) From the connection coefficients show that the low-velocity limit (23.67) of the geodesic equation is given by (23.68).

23.5.3 Uniformly Accelerated Observer

As an important example, consider an observer whose accelerated world line, written in some inertial (Lorentz) coordinate system $\{t, x, y, z\}$, is

$$t = (1/\kappa) \sinh(\kappa\tau), \quad x = y = 0, \quad z = (1/\kappa) \cosh(\kappa\tau). \quad (23.69)$$

Here τ is proper time along the world line, and κ is the magnitude of the observer's 4-acceleration, $\kappa = |\vec{a}|$ (which is constant along the world line); See Ex. 23.15, where the reader can derive the various claims made in this section.

This world line is depicted in Fig. 23.7 as a thick, solid hyperbola that asymptotes to the past light cone at early times and the future light cone at late times. The dots along the world line mark events that have proper times $\tau = -1.2, -0.9, -0.6, -0.3, 0, +0.3, +0.6, +0.9, +1.2$ (in units of $1/\kappa$). At each of these dots, the 3-plane orthogonal to the world line is shown as a dashed line (with the two dimensions out of the world line's 2-plane suppressed from the diagram). This 3-plane is labeled by its coordinate time $x^{\hat{0}}$, which is equal to the proper time of the dot. The basis vector $\vec{e}_{\hat{1}}$ is chosen to point along the observer's 4-acceleration, so $\vec{a} = \kappa \vec{e}_{\hat{1}}$. The coordinate $x^{\hat{1}}$ measures proper distance along the straight line that starts out tangent to $\vec{e}_{\hat{1}}$. The other two basis vectors $\vec{e}_{\hat{2}}$ and $\vec{e}_{\hat{3}}$ point out of the plane of the figure, and are parallel transported along the world line, $\nabla_{\vec{U}} \vec{e}_{\hat{2}} = \nabla_{\vec{U}} \vec{e}_{\hat{3}} = 0$; and $x^{\hat{2}}$ and $x^{\hat{3}}$ are measured along straight lines, in the orthogonal 3-plane, that start out tangent to these vectors. This construction implies that the resulting proper reference frame has vanishing rotation, $\vec{\Omega} = 0$ (Ex. 23.15), and that $x^{\hat{2}} = y$, $x^{\hat{3}} = z$, where y and z are coordinates in the system we used to define the world line [Eq. (23.69)].

Usually, in constructing an observer's proper reference frame, one confines attention to the immediate vicinity of her world line. However, in this special case it is instructive to




Fig. 23.7: The proper reference frame of a uniformly accelerated observer. All lengths and times are measured in units of $1/\kappa$. We show only two dimensions of the reference frame — those in the 2-plane of the observer’s curved world line.

extend the construction (the orthogonal 3-planes and their resulting space-time coordinates) outward arbitrarily far. By doing so, we discover that the 3-planes all cross at location $x^{\hat{1}} = -1/\kappa$, which means the coordinate system $\{x^{\hat{\alpha}}\}$ goes singular there. This singularity shows up in a vanishing $g_{\hat{0}\hat{0}}(x^{\hat{1}} = 1/\kappa)$ for the spacetime metric, written in that coordinate system:

$$ds^2 = -(1 + \kappa x^{\hat{1}})^2 (dx^{\hat{0}})^2 + (dx^{\hat{1}})^2 + (dx^{\hat{2}})^2 + (dx^{\hat{3}})^2 . \quad (23.70)$$

[Note that for $|x^{\hat{1}}| \ll 1/\kappa$ this metric agrees with the general proper-reference-frame metric (23.60b)]. From Fig. 23.7, it should be clear that this coordinate system can only cover, smoothly, one quadrant of Minkowski space-time: the quadrant $x > |t|$.

23.5.4 Rindler Coordinates for Minkowski Spacetime

The spacetime metric (23.70) in our observer’s proper reference frame resembles the metric in the vicinity of a black hole, as expressed in coordinates of observers who accelerate so as to avoid falling into the hole. In preparation for discussing this in Chap. 25, we shall shift the origin of our proper-reference-frame coordinates to the singular point, and rename them. Specifically, we introduce so-called *Rindler coordinates*²

$$t' = x^{\hat{0}} , \quad x' = x^{\hat{1}} + 1/\kappa , \quad y' = x^{\hat{2}} , \quad z' = x^{\hat{3}} . \quad (23.71)$$

²Named for Wolfgang Rindler.

It turns out (Ex. 23.15) that these coordinates are related to the Lorentz coordinates that we began with, in Eq. (23.69), by

$$t = x' \sinh(\kappa t'), \quad x = x' \cosh(\kappa t'), \quad y = y', \quad z = z'. \quad (23.72)$$

The metric in this Rindler coordinate system, of course, is the same as (23.70) with displacement of the origin:

$$ds^2 = -(\kappa x')^2 dt'^2 + dx'^2 + dy'^2 + dz'^2. \quad (23.73)$$

The world lines of constant $\{x', y', z'\}$ have uniform acceleration, $\vec{a} = (1/x')\vec{e}_{x'}$. Thus, we can think of these coordinates as the reference frame of a *family* of uniformly accelerated observers, each of whom accelerates away from their *horizon* $x' = 0$ with acceleration equal to $1/(\text{her distance } x' \text{ above the horizon})$. (We use the name “horizon” for $x' = 0$ because it represents the edge of the region of spacetime that these observers are able to observe.) The local 3-planes orthogonal to these observers’ world lines all mesh to form global 3-planes of constant t' . This is a major factor in making the metric (23.73) so simple.

EXERCISES

Exercise 23.15 *Derivation: Uniformly Accelerated Observer and Rinder Coordinates*

In this exercise you will derive the various claims made in Secs. 23.5.3 and 23.5.4

- Show that the parameter τ along the world line (23.69) is proper time, and that the 4-acceleration has magnitude $|\vec{a}| = 1/\kappa$.
- Show that the unit vectors $\vec{e}_{\hat{j}}$ introduced in Sec. 23.5.3 all obey the Fermi-Walker transport law (23.62) and therefore, by virtue of Eq. (23.61b), the proper reference frame built from them has vanishing rotation, $\vec{\Omega} = 0$.
- Show that the coordinates $x^{\hat{2}}$ and $x^{\hat{3}}$ introduced in Sec. 23.5.3 are equal to the y and z coordinates of the inertial frame used to define the observer’s world line [Eq. (23.69)].
- Show that the proper-reference-frame coordinates constructed in Sec. 23.5.3 are related to the original $\{t, x, y, z\}$ coordinates by

$$t = (x^{\hat{1}} + 1/\kappa) \sinh(\kappa x^{\hat{0}}), \quad x = (x^{\hat{1}} + 1/\kappa) \cosh(\kappa x^{\hat{0}}), \quad y = x^{\hat{2}}, \quad z = x^{\hat{3}}; \quad (23.74)$$

and from this, deduce the form (23.70) of the Minkowskii spacetime metric in the observer’s proper reference frame.

- Show that, when converted to Rindler coordinates by moving the spatial origin, the coordinate transformation (23.74) becomes (23.72), and the metric (23.70) becomes (23.73).
- Show that observers at rest in the Rindler coordinate system, i.e. who move along world lines of constant $\{x', y', z'\}$, have 4-acceleration $\vec{a} = (1/x')\vec{e}_{x'}$.

Exercise 23.16 *Example: Rigidly Rotating Disk*

Consider a thin disk, at $z = 0$ in a Lorentz reference frame, with radius R . The disk rotates rigidly with angular velocity Ω . In the early years of special relativity there was much confusion over the geometry of the disk: In the inertial frame it has physical radius (proper distance from center to edge) R and physical circumference $\mathcal{C} = 2\pi R$. But Lorentz contraction dictates that, as measured on the disk, the circumference should be $\sqrt{1 - v^2} \mathcal{C}$ (with $v = \Omega R$), and the physical radius, R , should be unchanged. This seemed weird. How could an obviously flat disk in flat spacetime have a curved, non-Euclidean geometry, with physical circumference divided by physical radius smaller than 2π ? In this exercise you will explore this issue.

- (a) Consider a family of observers who ride on the edge of the disk. Construct a circular curve, orthogonal to their world lines, that travels around the disk (at $\sqrt{x^2 + y^2} = R$). This curve can be thought of as lying in a 3-surface of constant time x^0 of the observers' proper reference frames. Show that it spirals upward in a Lorentz-frame spacetime diagram, so it cannot close on itself after traveling around the disk. This means that the 3-planes, orthogonal to the observers' world lines at the edge of the disk, cannot mesh globally to form global 3-planes (by contrast with the case of the uniformly accelerated observers in Sec. 23.5.4 and Ex. 23.15).
- (b) Next, consider a 2-dimensional family of observers who ride on the surface of the rotating disk. Show that at each radius $\sqrt{x^2 + y^2} = \text{constant}$, the constant-radius curve that is orthogonal to their world lines spirals upward in spacetime with a different slope. Show this means that even locally, the 3-planes orthogonal to each of their world lines cannot mesh to form larger 3-planes — and therefore, there does not reside, in spacetime, any 3-surface orthogonal to these observers' world lines. There is no 3-surface that has the claimed non-Euclidean geometry.

Bibliographic Note

For a very readable presentation of most of this chapter's material, from much the same point of view, see Chap. 20 of Hartle (2003). For an equally elementary introduction from a somewhat different viewpoint, see Chaps. 1–4 of Schutz (1980). A far more detailed and somewhat more sophisticated introduction, largely but not entirely from our viewpoint, will be found in Chaps. 1–6 of Misner, Thorne and Wheeler (1973). More sophisticated treatments from rather different viewpoints than ours are given in Chaps. 1 and 2 and Sec. 3.1 of Wald (1984), and in Chaps. 1 and 2 of Carroll (2004). A treasure trove of exercises on this material, with solutions, will be found in Chaps. 6, 7, and 8 of Lightman, Press, Price and Teukolsky (1975).

Box 23.4
Important Concepts in Chapter 23

- Most important concepts from Chap. 2
 - Principle of Relativity, Sec. 23.2.1
 - Metric defined in terms of interval, Sec. 23.2.1
 - Inertial frames, Sec. 23.2.2
 - Interval and spacetime diagrams, Sec. 23.2.3
- Differential geometry in general bases, Sec. 23.3
 - Dual bases, $\{\vec{e}_\alpha\}$, $\{\vec{e}^\mu\}$ with $\vec{e}_\alpha \cdot \vec{e}^\mu = \delta_\alpha^\mu$, Sec. 23.3.1
 - Covariant, contravariant and mixed components of a tensor, Sec. 23.3.1
 - Changes of bases and corresponding transformation of components of tensors, Sec. 23.3.1
 - Coordinate bases, Sec. 23.3.1
 - Orthonormal bases, Sec. 23.2.2
 - Vector as a differential operator (directional derivative), Sec. 23.3.2
 - Tangent space, Sec. 23.3.2
 - Commutator of vector fields, Sec. 23.3.2
 - Parallel transport of vectors, Sec. 23.3.3
 - Connection coefficients, how to compute them, and how to use them in computing components of the gradients of tensor fields, Sec. 23.3.3
 - Christoffel symbols (connection coefficients in a coordinate basis), Sec. 23.3.3
 - Levi-Civita tensor and its components, Sec. 23.3.4
 - Volume elements for integration, Sec. 23.3.4
- Stress-energy tensor, Sec. 23.4
 - For perfect fluid, Sec. 23.4
 - For point particle, Sec. 23.4
 - For electromagnetic field, Ex. 2.28 of Chap 2.
- Proper reference frame of an accelerated observer and metric in it, Sec. 23.5
 - Inertial guidance system and gyroscopes, Box 23.3
 - Fermi-Walker Transport law for inertial-guidance gyroscope, Eq. (23.62)
 - Metric in proper reference frame: Eq. (23.60)
 - Geodesic law of motion, Sec. 23.5.2
 - Uniformly accelerated observers, Sec. 23.5.3
 - Rindler coordinates, Sec. 23.5.4

Bibliography

Carroll, S. M., 2004. *Spacetime and Geometry: An Introduction to General Relativity*, San Francisco: Addison Wesley.

Feynman, R. P., 1966. *The Character of Physical Law*, Cambridge MA: MIT Press.

Hartle, J. B., 2003. *Gravity: An Introduction to Einstein's General Relativity*, San Francisco: Addison-Wesley. For auxiliary material, including a Mathematica program to compute connection coefficients, see <http://web.physics.ucsb.edu/~gravitybook/>.

Lightman, A. P., Press, W. H., Price, R. H. & Teukolsky, S. A. 1975. *Problem Book In Relativity and Gravitation*, Princeton: Princeton University Press.

MTW: Misner, Charles W., Thorne, Kip S., and Wheeler, John A. 1973. *Gravitation*, San Francisco: W. H. Freeman.

Ni, W.-T., and Zimmermann, M., 1978. "Inertial and gravitational effects in the proper reference frame of an accelerated, rotating observer," *Physical Review D*, **17**, 1473.

Pais, A. 1982. 'Subtle is the Lord...' *The Science and Life of Albert Einstein*, Oxford U.K.: Oxford University Press.

Schutz, B. 1980. *Geometrical Methods of Mathematical Physics*, Cambridge: Cambridge University Press.

Wald, R. M. 1984. *General Relativity*, Chicago: University of Chicago Press.

Will, C. M. 1993. *Was Einstein Right*, New York: Basic Books.